



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C07D 487/22, C07C 211/22, 211/23,</b> <b>211/18, 215/14, 257/18, A61P 35/00,</b> <b>A61K 31/155, 31/13, 31/40</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/66587</b>  <b>(43) International Publication Date:</b> 9 November 2000 (09.11.00)
<b>(21) International Application Number:</b> PCT/US00/11591  <b>(22) International Filing Date:</b> 27 April 2000 (27.04.00)  <b>(30) Priority Data:</b> 60/131,779                      30 April 1999 (30.04.99)                      US  <b>(71) Applicant (for all designated States except US):</b> SLIL BIOMEDICAL CORPORATION [US/US]; 535 Science Drive, Madison, WI 53711-1066 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> FRYDMAN, Benjamin [AR/US]; 821 N. Holt Circle, Madison, WI 53719 (US). MARTON, Laurence, J. [US/US]; 5810 Tree Line Drive, Madison, WI 53711 (US). REDDY, Venodhar, K. [IN/US]; 4446 Hillcrest Drive #3, Madison, WI 53705 (US). VALASINAS, Aldonia [AR/US]; 821 N. Holt Circle, Madison, WI 53719 (US). BLOKHIN, Andrei, V. [RU/US]; 14 Dorchester Circle, Madison, WI 53719 (US). BASU, Hirak, S. [US/US]; 7226 Colony Drive, Madison, WI 53717 (US).  <b>(74) Agents:</b> CERPA, Robert, K. et al.; Morrison & Foerster LLP, 755 Page Mill Road, Palo Alto, CA 94304-1018 (US).	<b>(81) Designated States:</b> AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished          upon receipt of that report.</i>	
<b>(54) Title:</b> CONFORMATIONALLY RESTRICTED POLYAMINE ANALOGS AS DISEASE THERAPIES		
<b>(57) Abstract</b>  Novel conformationally restricted polyamine analogs are provided, as well as compositions comprising these novel polyamine analogs. Methods of using the novel polyamine analogs in treatment of diseases such as cancer are also provided. Also provided is a method of delivering these analogs specifically to tumor cells by covalently attaching polyamine analogs to porphyrin compounds, along with novel polyamine-porphyrin covalent conjugates.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakistan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## CONFORMATIONALLY RESTRICTED POLYAMINE ANALOGS AS DISEASE THERAPIES

### CROSS-REFERENCE TO RELATED APPLICATIONS

5           This application claims priority benefit of co-pending provisional patent application  
U.S. Serial No. 60/131,779, filed on April 30, 1999. The content of that application is  
hereby incorporated by reference herein in its entirety.

## STATEMENT OF RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH

Not applicable.

## TECHNICAL FIELD

15           This invention relates to conformationally restricted polyamine analogs, and to porphyrin-polyamine conjugates. This invention also relates to uses of these analogs or conjugates in various medicinal applications, including suppressing cell proliferation and treating various diseases, including cancer.

## BACKGROUND OF THE INVENTION

20 Natural polyamines, e.g., spermidine, norspermidine, homospermidine, 1,4-diaminobutane (putrescine), and spermine, are simple aliphatic amines produced in eukaryotic cells by a highly regulated metabolic apparatus. Polyamine levels and the activity of the polyamine biosynthetic apparatus tend to be high in dividing mammalian cells and low in quiescent cells. Populations of cells depleted of their polyamine content  
25 stop growing and may die. Janne et al. (1978) *A. Biochim. Biophys. Acta.* 473:241 and Pegg et al. (1982) *Am. J. Cell. Physiol.* 243:212-221. Polyamines are reviewed in Morgan (1998) *Methods. Mol. Biol.* 79:3-30.

Several lines of evidence indicate that polyamines, particularly spermidine, are required for cell proliferation: (i) they are found in greater amounts in growing than in non-growing tissues; (ii) prokaryotic and eukaryotic mutants deficient in polyamine biosynthesis are auxotrophic for polyamines; and (iii) inhibitors specific for polyamine biosynthesis also inhibit cell growth. Despite this evidence, the precise biological role of

polyamines in cell proliferation is uncertain. It has been suggested that polyamines, by virtue of their charged nature under physiological conditions and their conformational flexibility, might serve to stabilize macromolecules, such as nucleic acids, by anion neutralization. Hafner et al. (1979) *J. Biol. Chem.* 254:12419; Pohjatipelto et al. (1981) *Nature* 293:475; Mamont et al. (1978) *Biochem. Biophys. Res. Commun.* 81:58; Bloomfield et al. (1981) in *Polyamines in Biology and Medicine*, Morris et al., Eds., Dekker, New York, pp. 183-205.

A treatment approach has been devised based on the observation that increases in the polyamine pool suppress polyamine biosynthesis. Porter et al. (1988) in *Advances in Enzyme Regulation*, Pergamon Press, pp. 57-79. This approach attempts to identify polyamine analogs which down-regulate polyamine biosynthesis, but which do not perform the polyamine functions required for cell growth. BESPM, a N-bis(ethyl) analog of spermine, has served as a model compound for this strategy. BESPM rapidly suppresses polyamine biosynthetic enzymes, depletes natural polyamine pools, and inhibits cell growth *in vitro*. Porter et al. (1987) *Cancer Res.* 47:2821-2825. In addition, BESPM suppresses polyamine uptake (Byers et al. (1990) *J. Physiol.* 142:460-467; and Kramer et al. (1993) *J. Cell. Physiol.* 115:399-407), and thus minimizes the ability of tumor cells to meet their polyamine requirement by taking them up from their environment. BESPM and related analogs also induce the polyamine metabolizing enzyme spermidine/spermine N<sup>1</sup>-acetyltransferase (SSAT) in certain human carcinoma cell lines.

BESPM and other polyamine analogs have been used, or proposed for use, in treating a large variety of diseases, including a number of different cancers. U.S. Patent No. 5,541,230. Polyamine analogs demonstrated, for example, potent antitumor activity against several melanoma cell lines and tumors *in vitro* (Porter et al. (1991) *Cancer Res.* 51:3715-3720; Shappell et al. (1992) *Anticancer Res.* 12:1083-1090) and *in vivo* using tumors growing as xenografts in athymic mice (Bernacki et al. (1992) *Cancer Res.* 52:2424-2430; Porter et al. (1993) *Cancer Res.* 53:581-586). Potent antitumor activity of bis-ethyl spermine analogs has also been demonstrated for pancreatic cancer cell lines *in vitro* (Chang et al. (1992) *Cancer Chemother. Pharmacol.* 30:183-188) and *in vivo* (Chang et al. (1992) *Cancer Chemother. Pharmacol.* 30:179-182). Polyamine analogs have also been suggested for use in treating brain tumor therapy. Redgate et al. (1995) *J. Neurooncol.* 25:167-79. In addition to being useful against cancers of the brain, pancreas, and skin,

polyamine analogs are also useful against cancers of the bladder, bone, breast, colon, digestive tract, lung and ovaries. Chang et al. (1993) *J. Urol.* 150:1293-7; Snyder et al. (1994) *Anticancer Res.* 14:347-56; Yuan et al. (1994) *Biochem. Pharmacol.* 47:1587-92; Davidson et al. (1993) *Cancer Res.* 53:2071-5; Berchtold et al. (1998) *J. Cell. Physiol.* 174:380-6; Porter et al. (1988) *Adv. Exp. Med. Biol.* 250:677-90; U.S. Patent Nos. 5,498,522 and 5,374,658. U.S. Patent No. 5,498,522 presents the use of spermidine/spermine N<sup>1</sup>-acetyltransferase as a prognostic indicator of the efficacy of a polyamine analog against a malignant tumor.

Polyamine analogs have been used to treat cancer of the prostate. Mi et al. (1988) *Prostate* 34:51-60. Polyamines are produced in large amounts by the prostate gland and are abundant in the seminal fluid. Herr et al. (1984) *Cancer* 53:1294-8. Polyamine analogs such as BE-4444, BE-373, and BE-333 are particularly effective in inhibiting prostate xenograft tumors in nude mice. Zagaja et al. (1998) *Cancer Chem. Pharm.* 41:505-512; Jeffers et al. (1997) *Cancer Chem. Pharm.* 40:172-179; Feuerstein et al. (1991) *J. Cell. Biochem.* 46:37-47; and Marton et al. (1995) *Ann. Rev. Pharm. Toxicol.* 35:55-91.

In addition to treating cancer, polyamines and their analogs have uses in treating a number of other diseases and in numerous other medicinal applications. Oxidized polyamines are believed to inhibit growth of parasites (Morgan et al. (1983) *Adv. Polyamine Res.* 4: 169-174; Morgan et al. (1986) *Biochem. J.* 236:97-101; and U.S. Patent No. 4,935,449) and suppress infectivity of selected strains of bacteria and fungi (Bachrach et al. (1971) *J. Gen. Virol.* 13:415-22; Nishimura et al. (1971) *Biochim. Biophys. Acta* 247:153-6; and U.S. Patent No. 5,744,453). Polyamines such as spermine and polyamine analogs are also anti-viral and some are anti-insecticidal. Bachrach et al. (1972) *Appl. Microbiol.* 23:232-5; Bachrach et al. (1971) *J. Gen. Virol.* 11:1-9; U.S. Patent Nos. 5,021,409; 5,606,053; 5,608,061; 5,612,478; and 5,681,837. In addition, oxidized polyamines, such as spermine dialdehyde, for example, can be used in treatment of tissue grafts and other organs for transplantation. U.S. Patent No. 5,374,658. Polyamine analogs can also be used to treat neurodegenerative diseases and neurotrauma such as stroke. U.S. Patent Nos. 5,646,188 and 5,677,349. Polyamine analogs have also been reported to be useful as anti-psoratic agents, and in the treatment of epilepsy, Alzheimer's disease, and multiple sclerosis, as described in U.S. Patent No. 5,646,188. Polyamine analogs are also useful in treating and preventing restenosis. U.S. Patent No. 5,516,807. Polyamine analogs

are also useful in treatment of gastric ulcers. Igarashi et al. (1990) *Biochem. Biophys. Res. Commun.* 172:715-20. In addition, polyamine derivatives including N-alkythio polyamine derivatives, polyamine thiols, and polyamine phenols are useful as radioprotective agents for normal tissues during radiotherapy. U.S. Patent Nos. 5,217,964; 5,354,782; and  
5 5,434,145.

Polyamines and their analogs can be administered alone or in conjunction with additional agents. For example, therapeutic polyamines can be administered along with 1,3-bis (2-chloroethyl)-1-nitrosourea. U.S. Patent No. 5,541,230. In treating cancer, polyamines can be co-administered with various cytotoxic agents, including antineoplastic  
10 vinca alkaloids, antibiotics, antimetabolites, and platinum coordination complexes. U.S. Patent No. 5,654,287.

In addition to the various aforementioned medicinal uses, polyamines and polyamine analogs have a variety of industrial uses, including derivativization of silica. U.S. Patent No. 5,763,388. Polyamines have also been used in conjunction with other  
15 clarification aids to treat wastewaters. U.S. Patent Nos. 5,413,719 and 5,707,532. The combination of aluminum chlorohydrate and a polyamine is an effective emulsion breaker for reverse (oil-in-water) emulsions, e.g. in a matrix comprising mostly oil as encountered in a crude oil desalter unit. U.S. Patent No. 5,607,574. Polyamines are also useful in deodorizing polysulfides. U.S. S.I.R. H1,633. Polyamines are also used in industrial dyes.  
20 U.S. Patent No. 5,672,202. Polyamines and hot water can also be used in manufacturing microcapsules. U.S. Patent No. 5,401,443. The antioxidative and metal-chelating effects of polyamines are reviewed in Lovaas (1997) *Adv. Pharmacol.* 38:119-149.

It would be advantageous to develop novel polyamine analogs for various uses, including disease treatment.

25 All references cited herein are hereby incorporated by reference in their entirety.

## SUMMARY OF THE INVENTION

The invention provides novel polyamine analogs, compositions comprising a polyamine analog, and methods using the analogs and compositions. In one embodiment,  
30 the polyamine analog is conformationally restricted.

In another embodiment, the polyamine analog is selected from among compounds of the formula:


E-NH-B-A-B-NH-B-A-B-NH-B-A-B-NH-B-A-B-NH-E


where A is independently selected from the group consisting of a single bond, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl; B is independently selected from the group consisting of: a single bond, C<sub>1</sub>-C<sub>6</sub> alkyl, and

5 C<sub>2</sub>-C<sub>6</sub> alkenyl; and E is independently selected from the group consisting of H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl;

with the proviso that either at least one A moiety is selected from the group consisting of C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl,

10 or at least one B moiety is selected from the group consisting of C<sub>2</sub>-C<sub>6</sub> alkenyl; and all salts and stereoisomers thereof. Specific embodiments of compounds of this type include






, and all salts and

stereoisomers thereof.

In another embodiment, the polyamine analog is selected from among the group of compounds of the formula:



wherein A is independently selected from the group consisting of a single bond, C<sub>6</sub>-C<sub>2</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub>

cycloalkenyl; B is independently selected from the group consisting of a single bond,

C<sub>1</sub>-C<sub>6</sub> alkyl, and C<sub>2</sub>-C<sub>6</sub> alkenyl; E is independently selected from the group consisting of

10 H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub>


cycloalkenyl; and x is an integer from 2 to 16; with the proviso that either at least one A

moiety is selected from the group consisting of C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub>


cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl, or at least one B moiety is selected

from the group consisting of C<sub>2</sub>-C<sub>6</sub> alkenyl; and all salts and stereoisomers thereof.

15 Specific embodiments of compounds of this type include




and



;

and all salts and stereoisomers thereof.



20 In another embodiment, the polyamine analog is selected from among the group of compounds of the formula






wherein A is independently selected from the group consisting of a single bond, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl; B is independently selected from the group consisting of a single bond, C<sub>1</sub>-C<sub>6</sub> alkyl, and C<sub>2</sub>-C<sub>6</sub> alkenyl; E is independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkanol, C<sub>3</sub>-C<sub>6</sub> cycloalkanol, and C<sub>3</sub>-C<sub>6</sub> hydroxyaryl, with the proviso that at least one E moiety be selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkanol, C<sub>3</sub>-C<sub>6</sub> cycloalkanol, and C<sub>3</sub>-C<sub>6</sub> hydroxyaryl; and x is an integer from 0 to 16; and all salts or stereoisomers thereof.

Specific embodiments of compounds of this type include




, and



;

and all salts and stereoisomers thereof.

In another embodiment, the polyamine analog is selected from among the group of compounds of the formula





wherein A is selected from the group consisting of C<sub>2</sub>-C<sub>6</sub> alkynyl; B is independently selected from the group consisting of: a single bond, C<sub>1</sub>-C<sub>6</sub> alkyl, and C<sub>2</sub>-C<sub>6</sub> alkenyl; D is independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloalkenyl, and C<sub>3</sub>-C<sub>6</sub> cycloaryl; and E is independently selected from the group consisting of H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub>

cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl; and all salts and stereoisomers thereof. Specific embodiments of compounds of this type include




5




10 and all salts and stereoisomers thereof.


In another embodiment, the polyamine analog is selected from among the group of compounds of the formula



15 wherein F is selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl; A is independently selected from the group consisting of: a single bond, C<sub>1</sub>-C<sub>6</sub> alkyl; C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl; B is independently selected from the group consisting of: a single bond, C<sub>1</sub>-C<sub>6</sub> alkyl, and C<sub>2</sub>-C<sub>6</sub> alkenyl; and E is independently selected from the group consisting of H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl; with the proviso that  
20 either at least one A moiety is selected from the group consisting of C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl, or at least one B moiety is selected from the group consisting of C<sub>2</sub>-C<sub>6</sub> alkenyl; and all salts and stereoisomers thereof. Specific embodiments of compounds of this type include




and



;



and all salts and stereoisomers thereof.

- 5 In another embodiment, the polyamine analog is selected from among the group of compounds of the formula




- 10 wherein A is independently selected from the group consisting of a single bond, C<sub>1</sub>-C<sub>6</sub> alkyl; C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl; B is independently selected from the group consisting of: a single bond, C<sub>1</sub>-C<sub>6</sub> alkyl, and C<sub>2</sub>-C<sub>6</sub> alkenyl; with the proviso that A and both B moieties are not all a single bond; and all salts and stereoisomers thereof. Specific embodiments of compounds of this type include

15




, and



and all salts and stereoisomers thereof.

In another embodiment, the polyamine analog is selected from among the group of compounds of the formula




wherein A is selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloalkenyl, and C<sub>3</sub>-C<sub>6</sub> cycloaryl; B is independently selected from the group consisting of: a single bond, C<sub>1</sub>-C<sub>6</sub> alkyl, and C<sub>2</sub>-C<sub>6</sub> alkenyl; D is independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloalkenyl, and C<sub>3</sub>-C<sub>6</sub> cycloaryl; and E is independently selected from the group consisting of H and C<sub>1</sub>-C<sub>6</sub> alkyl; with the proviso that either at least one A moiety is selected from the group consisting of C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl, or at least one B moiety is selected from the group consisting of C<sub>2</sub>-C<sub>6</sub> alkenyl; and the proviso that at least one E is selected from the group consisting of H or methyl; and all salts and stereoisomers thereof. A specific embodiment of compounds of this type includes



and all salts and stereoisomers thereof.


In another embodiment, the polyamine analog is selected from among the group of compounds of the formula



wherein A is *trans*-ethene (no stereochemical variation about the double bond being permitted, that is, *cis*-ethene is specifically excluded); B is independently selected from the

group consisting of: a single bond, C<sub>1</sub>-C<sub>6</sub> alkyl, and C<sub>2</sub>-C<sub>6</sub> alkenyl; D is independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl, with the proviso that D is not C<sub>3</sub> alkyl; and E is independently selected from the group consisting of H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl; and all salts thereof.

In another embodiment the invention also provides conjugates of polyamines to porphyrin compounds, where the polyamines are linked via a covalent bond to the porphyrin compound. Polyamines useful for this embodiment of the invention include, but are not limited to, the polyamines described in the previous embodiments, or the polyamines depicted in Table 1. The covalent bond can be an amide bond, an amine bond, or any other suitable covalent bond. The polyamine analogs can be bound to the porphyrin compound at positions including, but not limited to, any of the peripheral positions of the porphyrin macrocycle such as the  $\beta$ -pyrrole positions or the meso carbons of the macrocycle. Non-limiting examples of these compounds are given in Table 3. In one embodiment, the porphyrin-polyamine compounds are of the formula



where at least one of J<sub>1</sub>-J<sub>8</sub> is independently selected from the group consisting of -K<sub>1</sub>-G-L-(N(P)-A)<sub>n</sub>-K<sub>2</sub>; where K<sub>1</sub> is independently selected from the group consisting of C<sub>1</sub>-C<sub>8</sub> alkyl and where the valence to the left of K<sub>1</sub> attaches to the porphyrin ring; where G is -O-, -(C=O)-, -C(=O)-O-, -O-(C=O)-, -O-(C=O)-O-, -O-(C=O)-N-, -N-(C=O)-O-, or a

nonentity; where L is C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>3</sub>-C<sub>8</sub> cycloaryl, C<sub>1</sub>-C<sub>8</sub> alkoxy, C<sub>1</sub>-C<sub>8</sub> alkyl-C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>1</sub>-C<sub>8</sub> alkyl-C<sub>3</sub>-C<sub>8</sub> cycloaryl, C<sub>1</sub>-C<sub>8</sub> alkoxy-C<sub>3</sub>-C<sub>8</sub> cycloaryl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl-C<sub>3</sub>-C<sub>8</sub> cycloaryl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl-C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>3</sub>-C<sub>8</sub> cycloaryl-C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>3</sub>-C<sub>8</sub> cycloaryl-C<sub>1</sub>-C<sub>8</sub> alkoxy, C<sub>3</sub>-C<sub>8</sub> cycloaryl-C<sub>3</sub>-C<sub>8</sub> cycloalkyl, or a nonentity; each A is independently selected from the group consisting of C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>2</sub>-C<sub>8</sub> alkenyl, C<sub>2</sub>-C<sub>8</sub> alkynyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>3</sub>-C<sub>8</sub> cycloaryl, C<sub>3</sub>-C<sub>8</sub> cycloalkenyl, and C<sub>3</sub>-C<sub>8</sub> cycloalkynyl; P is selected from the group consisting of H and C<sub>1</sub>-C<sub>8</sub> alkyl; n is an integer from 2 to 8; and K<sub>2</sub> is independently selected from the group consisting of H, C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>2</sub>-C<sub>8</sub> alkenyl, C<sub>2</sub>-C<sub>8</sub> alkynyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>3</sub>-C<sub>8</sub> cycloaryl, C<sub>3</sub>-C<sub>8</sub> cycloalkenyl, C<sub>3</sub>-C<sub>8</sub> cycloalkynyl, C<sub>1</sub>-C<sub>8</sub> alkanol, C<sub>3</sub>-C<sub>8</sub> cycloalkanol, and C<sub>3</sub>-C<sub>8</sub> hydroxyaryl; where the remainder of the groups J<sub>1</sub>-J<sub>8</sub> are each independently selected from the group consisting of H, K<sub>3</sub>, or K<sub>4</sub>-COOH, where K<sub>3</sub> is independently selected from the group consisting of C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>2</sub>-C<sub>8</sub> alkenyl, C<sub>2</sub>-C<sub>8</sub> alkynyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>3</sub>-C<sub>8</sub> cycloaryl, C<sub>3</sub>-C<sub>8</sub> cycloalkenyl, C<sub>3</sub>-C<sub>8</sub> cycloalkynyl, C<sub>1</sub>-C<sub>8</sub> alkanol, C<sub>3</sub>-C<sub>8</sub> cycloalkanol, and C<sub>3</sub>-C<sub>8</sub> hydroxyaryl; and K<sub>4</sub> is independently selected from the group consisting of C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>2</sub>-C<sub>8</sub> alkenyl, C<sub>2</sub>-C<sub>8</sub> alkynyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>3</sub>-C<sub>8</sub> cycloaryl, C<sub>3</sub>-C<sub>8</sub> cycloalkenyl, and C<sub>3</sub>-C<sub>8</sub> cycloalkynyl; and any salt or stereoisomer thereof.

Additional embodiments include those where K<sub>1</sub> and each A are independently selected from the group consisting of C<sub>1</sub>-C<sub>8</sub> alkyl, K<sub>2</sub> is independently selected from the group consisting of H and C<sub>1</sub>-C<sub>8</sub> alkyl, and the remainder of the groups J<sub>1</sub>-J<sub>8</sub> are each independently selected from the group consisting of H, K<sub>3</sub>, or K<sub>4</sub>-COOH, where K<sub>3</sub> and K<sub>4</sub> are independently selected from the group consisting of C<sub>1</sub>-C<sub>8</sub> alkyl. In additional embodiments, K<sub>1</sub> is -CH<sub>2</sub>-CH<sub>2</sub>- or -CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-. In additional embodiments, n is 4. In yet additional embodiments, J<sub>3</sub>, J<sub>4</sub>, J<sub>7</sub> and J<sub>8</sub> of the formula depicted above are independently selected from the group consisting of C<sub>1</sub>-C<sub>3</sub> alkyl; J<sub>5</sub> and J<sub>6</sub> are independently selected from the group consisting of C<sub>1</sub>-C<sub>3</sub> alkyl and C<sub>1</sub>-C<sub>3</sub> alkyl-COOH; and J<sub>1</sub> and J<sub>2</sub> are independently selected from the group consisting of C<sub>1</sub>-C<sub>3</sub> alkyl-G-N(P<sub>1</sub>)-A-(NH-A)<sub>n1</sub>-K<sub>5</sub>; where G is -(C=O)- or a nonentity; P<sub>1</sub> is H or C<sub>1</sub>-C<sub>3</sub> alkyl; each A is independently selected from the group consisting of C<sub>1</sub>-C<sub>8</sub> alkyl; n<sub>1</sub> is 3 or 4; and K<sub>5</sub> is independently selected from the group consisting of H and C<sub>1</sub>-C<sub>8</sub> alkyl.

The invention also includes methods of treating an indication in an individual comprising the step of administering to the individual a therapeutic amount of a porphyrin-polyamine conjugate of the invention.

The invention also provides methods of treating an indication comprising the step of administering to the individual an effective amount of a composition comprising a polyamine analog, preferably a conformationally restricted polyamine analog.

In one method, the indication is cancer. In various embodiments, the cancer affects cells of the bladder, blood, brain, breast, colon, digestive tract, lung, ovaries, pancreas, prostate gland, or skin. In other embodiments, the indication can also include, but is not limited to, Alzheimer's disease, epilepsy, multiple sclerosis, problems associated with tissue grafts and organ transplants, psoriasis, restenosis, stomach ulcers, or tissue overgrowth after surgery. In other embodiments, the indication is an infection or infestation of parasites, bacteria, fungi or insects. The polyamine analog or porphyrin-polyamine conjugate is selected from the foregoing groups of compounds.

## BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a graph depicting the *in vitro* effect of increasing concentrations of BE-4444 (○), SL-11121 (●), SL-11122 (□), SL-11123 (■), SL-11126 (Δ), SL-11127 (▲), SL-11128 (◇), SL-11129 (◆), SL-11130 (▽), SL-11133 (▼) on the survival of cultured human prostate cancer cells PC3.

ED<sub>50</sub> of BE-4444 = 0.6 μM, SL-11121 = 0.52 μM, SL-11122 >31.25 μM,

SL-11123 >31.25 μM, SL-11126 = 0.2 μM SL-11127 >31.25 μM,

SL-11128 = 0.5 μM, SL-11129 = 1.7 μM, SL-11130 >31.25 μM, and

SL-11133 >31.25 μM.

Fig. 2 is a graph depicting the *in vitro* effect of increasing concentrations of BE-4444 (○), SL-11121 (●), SL-11122 (□), SL-11123 (■), SL-11126 (Δ), SL-11127 (▲), SL-11128 (◇), SL-11129 (◆), SL-11130 (▽), and SL-11133 (▼) on the survival of cultured human prostate cancer cells DU145.

ED<sub>50</sub> of BE-4444 = 0.07 μM, SL-11121 = 0.08 μM, SL-11122 = 0.08 μM,

SL-11123 = 0.51 μM, SL-11126 = 0.51 μM SL-11127 0.22 μM,

SL-11128 = 0.14  $\mu$ M, SL-11129 = 0.32  $\mu$ M, SL-11130 = 0.43  $\mu$ M, and  
SL-11133 = 0.34  $\mu$ M.

Fig. 3 is a graph depicting the *in vitro* effect of increasing concentrations of BE-4444 (○), SL-11121 (●), SL-11122 (□), SL-11123 (■), SL-11126 (Δ), SL-11127 (▲), SL-11128 (◇), SL-11129 (◆), SL-11130 (▽), and SL-11133 (▼) on the survival of cultured human prostate cancer cells DUPRO.

ED<sub>50</sub> of BE-4444 = 0.2  $\mu$ M, SL-11121 = 0.4  $\mu$ M, SL-11122 = 0.56  $\mu$ M,  
SL-11123 >31.25  $\mu$ M, SL-11126 = 1.1  $\mu$ M, SL-11127 1.3  $\mu$ M,  
SL-11128 = 1.28  $\mu$ M, SL-11129 >31.25  $\mu$ M, SL-11130 >31.25  $\mu$ M, and  
SL-11133 = 31.25  $\mu$ M.

Fig. 4 is a graph depicting the *in vitro* effect of increasing concentrations of BE-4444 (○), SL-11121 (●), SL-11126 (Δ), SL-11128 (◇), on the survival of cultured human prostate cancer cells LNCAP.

ED<sub>50</sub> of BE-4444 = 0.14  $\mu$ M, SL-11121 = 0.14  $\mu$ M, SL-11126 = 0.55  $\mu$ M and  
SL-11128 = 0.3  $\mu$ M.

Fig. 5 is a graph depicting the *in vitro* effect of increasing concentrations of BE-4444 (○), SL-11121 (●), SL-11122 (□), SL-11123 (■), SL-11126 (Δ), SL-11127 (▲), and SL-11128 (◇) on the survival of cultured human colon cancer cells HT29.

ED<sub>50</sub> of BE-4444 = 0.5  $\mu$ M, SL-11121 = 0.8  $\mu$ M, SL-11122 = 0.8  $\mu$ M,  
SL-11123 = 10.42  $\mu$ M, SL-11126 = 1.5  $\mu$ M, SL-11127 = 2.91  $\mu$ M, and  
SL-11128 = 1.35  $\mu$ M.

Fig. 6 is a graph depicting the *in vitro* effect of increasing concentrations of BE-4444 (○), SL-11121 (●), SL-11122 (□), SL-11123 (■), and SL-11126 (Δ) on the survival of cultured human lung cancer cells A549.

ED<sub>50</sub> of BE-4444 >31.25  $\mu$ M, SL-11121 >31.25  $\mu$ M, SL-11122 >31.25  $\mu$ M,  
SL-11123 >31.25  $\mu$ M, and SL-11126 >31.25  $\mu$ M.

Fig. 7 is a graph depicting the *in vitro* effect of increasing concentrations of BE-4444 (○), SL-11121 (●), SL-11122 (□), SL-11123 (■), and SL-11126 (Δ) on the survival of cultured human breast cancer cells MCF7.

ED<sub>50</sub> of BE-4444 >31.25  $\mu$ M, SL-11121 = 17.0  $\mu$ M, SL-11122 >31.25  $\mu$ M,  
SL-11123 >31.25  $\mu$ M, and SL-11126 = 0.7  $\mu$ M.



Fig. 8 is a graph depicting the *in vitro* effect of increasing concentrations of SL-11105 (●), SL-11124 (○), SL-11132 (■), and BE-333 (□) on the survival of cultured human prostate cancer cells PC3.

ED<sub>50</sub> of SL-11105 >31.25 μM, SL-11124 >31.25 μM, SL-11132 >31.25 μM  
5 and BE-333 = 0.34 μM.

Fig. 9 is a graph depicting the *in vitro* effect of increasing concentrations of SL-11105 (●), SL-11124 (○), SL-11132 (■), and BE-333 (□) on the survival of cultured human prostate cancer cells DU145.

ED<sub>50</sub> of SL-11105 = 1.6 μM, SL-11124 >31.25 μM, SL-11132 = 0.015 μM  
10 and BE-333 = 0.12 μM.

Fig. 10 is a graph depicting the *in vitro* effect of increasing concentrations of SL-11105 (●), SL-11124 (○), SL-11132 (■), and BE-333 (□) on the survival of cultured human prostate cancer cells DUPRO.

ED<sub>50</sub> of SL-11105 = 0.43 μM, SL-11124 >31.25 μM, SL-11132 >31.25 μM  
15 and BE-333 = 0.9 μM.

Fig. 11 is a graph depicting the *in vitro* effect of increasing concentrations of SL-11105 (●), SL-11124 (○), and BE-333 (□) on the survival of cultured human colon cancer cells HT29.

ED<sub>50</sub> of SL-11105 = 25.2 μM, SL-11124 >31.25 μM, and BE-333 = 0.3 μM.

Fig. 12 is a graph depicting the *in vitro* effect of increasing concentrations of SL-11105 (●), SL-11124 (○), and BE-333 (□) on the survival of cultured human lung cancer cells A549.

ED<sub>50</sub> of SL-11105 = 0.43 μM, SL-11124 >31.25 μM, and BE-333 = 0.3 μM.

Fig. 13 is a graph depicting the *in vitro* effect of increasing concentrations of SL-11105 (●), SL-11124 (○), and BE-333 (□) on the survival of cultured human breast cancer cells MCF7.

ED<sub>50</sub> of SL-11105 >31.25 μM, SL-11124 >31.25 μM, and BE-333 = 3.7 μM.

Fig. 14 is a graph depicting the *in vitro* effect of increasing concentrations of SL-11105 (●) and BE-333 (□) on the survival of cultured human brain tumor cells U251  
30 MG NCI.

ED<sub>50</sub> of SL-11105 = 25.9 μM, and BE-333 = 0.23 μM.

Fig. 15A is a graph depicting the *in vitro* effect of increasing concentrations of SL-11093 (●), SL-11098 (○), SL-11099 (■), SL-11100 (□), SL-11101 (▲), SL-11102 (Δ), and BE-444 (◆) on the survival of cultured human prostate cancer cells PC3.

ED<sub>50</sub> of SL-11093 = 1.6 μM, SL-11098 = 1.4 μM, SL-11099 = 2.5 μM,  
 5 SL-11100 = 4.7 μM, SL-11101 = 7.7 μM, SL-11102 >31.25 μM and  
 BE-444 = 0.7 μM.

Fig. 15B is a graph depicting the *in vitro* effect of increasing concentrations of SL-11103 (●), SL-11104 (○), SL-11108 (■), SL-11114 (□), SL-11118 (▲), SL-11119 (Δ), and BE-444 (◆) on the survival of cultured human prostate cancer cells PC3.

10 ED<sub>50</sub> of SL-11103 >31.25 μM, SL-11104 >31.25 μM, SL-11108 = 2.2 μM,  
 SL-11114 = 0.7 μM, SL-11118 = 1.65 μM, SL-11119 >31.25 μM and  
 BE-444 = 0.7 μM

Fig. 16A is a graph depicting the *in vitro* effect of increasing concentrations of SL-11093 (●), SL-11098 (○), SL-11099 (■), SL-11100 (□), SL-11101 (▲), SL-11102 (Δ), and  
 15 BE-444 (◆) on the survival of cultured human prostate cancer cells DU145.

ED<sub>50</sub> of SL-11093 = 0.016 μM, SL-11098 = 0.02 μM, SL-11099 = 0.014 μM,  
 SL-11100 = 0.021 μM, SL-11101 = 0.22 μM, SL-11102 = 0.03 μM and  
 BE-444 = 0.03 μM.

Fig. 16B is a graph depicting the *in vitro* effect of increasing concentrations of SL-11103 (●), SL-11104 (○), SL-11108 (■), SL-11114 (□), SL-11118 (▲), SL-11119 (Δ), and  
 20 BE-444 (◆) on the survival of cultured human prostate cancer cells DU145.

ED<sub>50</sub> of SL-11103 = 2.8 μM, SL-11104 = 9.4 μM, SL-11108 = 0.13 μM,  
 SL-11114 = 0.13 μM, SL-11118 = 0.05 μM, SL-11119 0.08 μM and  
 BE-444 = 0.03 μM.

25 Fig. 17A is a graph depicting the *in vitro* effect of increasing concentrations of SL-11099 (■), SL-11100 (□), SL-11101 (▲), SL-11102 (Δ), and BE-444 (◆) on the survival of cultured human prostate cancer cells DUPRO.

ED<sub>50</sub> of = SL-11099 = 0.08 μM, SL-11100 = 0.3 μM, SL-11101 = 0.85 μM,  
 SL-11102 = 0.15 μM and BE-444 = 0.2 μM.

**Fig. 17B** is a graph depicting the *in vitro* effect of increasing concentrations of SL-11108 (■), SL-11114 (□), SL-11118 (▲), SL-11119 (Δ), and BE-444 (◆) on the survival of cultured human prostate cancer cells DUPRO.

ED<sub>50</sub> of SL-11108 = 0.98 μM, SL-11114 = 0.64 μM, SL-11118 = 0.25 μM,

5 SL-11119 = 0.44 μM and BE-444 = 0.2 μM.

**Fig. 18A** is a graph depicting the *in vitro* effect of increasing concentrations of SL-11093 (●), SL-11098 (○), SL-11099 (■), SL-11100 (□), and BE-444 (◆) on the survival of cultured human prostate cancer cells LNCAP.

ED<sub>50</sub> of SL-11093 = 0.21 μM, SL-11098 = 0.17 μM, SL-11099 = 0.21 μM,

10 SL-11100 = 0.7 μM, and BE-444 = 0.1 μM.

**Fig. 18B** is a graph depicting the *in vitro* effect of increasing concentrations of SL-11108 (■), SL-11114 (□), SL-11118 (▲), and BE-444 (◆) on the survival of cultured human prostate cancer cells LNCAP.

ED<sub>50</sub> of SL-11108 = 7.7 μM, SL-11114 = 3.0 μM, SL-11118 = 0.21 μM,

15 and BE-444 = 0.1 μM.

**Fig. 19A** is a graph depicting the *in vitro* effect of increasing concentrations of SL-11093 (●), SL-11098 (○), SL-11099 (■), SL-11100 (□), SL-11101 (▲), SL-11102 (Δ), and BE-444 (◆) on the survival of cultured human colon cancer cells HT29.

ED<sub>50</sub> of SL-11093 = 0.4 μM, SL-11098 = 0.4 μM, SL-11099 = 1.0 μM,

20 SL-11100 = 2.0 μM, SL-11101 = 5.2 μM. SL-11102 = 0.73 μM and

BE-444 = 0.93 μM.

**Fig. 19B** is a graph depicting the *in vitro* effect of increasing concentrations of SL-11103 (●), SL-11104 (○), SL-11108 (■), SL-11114 (□), SL-11118 (▲), SL-11119 (Δ), and BE-444 (◆) on the survival of cultured human colon cancer cells HT29.

25 ED<sub>50</sub> of SL-11103 = 29.4 μM, SL-11104 = 25.8 μM, SL-11108 = 2.0 μM,

SL-11114 = 3.6 μM, SL-11118 = 0.98 μM, SL-11119 = 0.97 μM and

BE-444 = 0.93 μM.

**Fig. 20A** is a graph depicting the *in vitro* effect of increasing concentrations of SL-11093 (●), SL-11098 (○), SL-11099 (■), SL-11100 (□), SL-11101 (▲), SL-11102 (Δ), and BE-444 (◆) on the survival of cultured human lung cancer cells A549.

30

ED<sub>50</sub> of SL-11093 = 0.26 μM, SL-11098 = 0.29 μM, SL-11099 = 0.51 μM,

SL-11100 = 0.65  $\mu$ M, SL-11101 = 2.2  $\mu$ M, SL-11102 = 0.15  $\mu$ M and  
BE-444 = 0.15  $\mu$ M.

Fig. 20B is a graph depicting the *in vitro* effect of increasing concentrations of SL-  
11103 (●), SL-11104 (○), SL-11108 (■), SL-11114 (□), SL-11118 (▲), and BE-444 (◆) on  
5 the survival of cultured human lung cancer cells A549.

ED<sub>50</sub> of SL-11103 = 12.4  $\mu$ M, SL-11104 >31.25  $\mu$ M, SL-11108 >31.25  $\mu$ M,  
SL-11114 >31.25  $\mu$ M, SL-11118 = 0.214  $\mu$ M and BE-444 = 0.15  $\mu$ M.

Fig. 21A is a graph depicting the *in vitro* effect of increasing concentrations of SL-  
11093 (●), SL-11098 (○), SL-11099 (■), SL-11100 (□), SL-11101 (▲), SL-11102 (Δ), and  
10 BE-444 (◆) on the survival of cultured human breast cancer cells MCF7.

ED<sub>50</sub> of SL-11093 = 0.66  $\mu$ M, SL-11098 >31.25  $\mu$ M, SL-11099 = 26.3  $\mu$ M,  
SL-11100 >31.25  $\mu$ M, SL-11101 >31.25  $\mu$ M SL-11102 >31.25  $\mu$ M and  
BE-444 >31.25  $\mu$ M.

Fig. 21B is a graph depicting the *in vitro* effect of increasing concentrations of SL-  
11103 (●), SL-11104 (○), SL-11108 (■), and BE-444 (◆) on the survival of cultured  
15 human breast cancer cells MCF7.

ED<sub>50</sub> of SL-11103 >31.25  $\mu$ M, SL-11104 >31.25  $\mu$ M, SL-11108 >31.25  $\mu$ M,  
and BE-444 >31.25  $\mu$ M.

Fig. 22A is a graph depicting the *in vitro* effect of increasing concentrations of SL-  
11093 (●), SL-11098 (○), SL-11099 (■), SL-11100 (□), SL-11101 (▲), SL-11102 (Δ), and  
20 BE-444 (◆) on the survival of cultured human brain tumor cells U251 MG NCI.

ED<sub>50</sub> of SL-11093 = 0.07  $\mu$ M, SL-11098 = 0.1  $\mu$ M, SL-11099 = 0.11  $\mu$ M,  
SL-11100 = 0.22  $\mu$ M, SL-11101 = 1.7  $\mu$ M, SL-11102 = 0.15  $\mu$ M and  
BE-444 = 0.2  $\mu$ M.

Fig. 22B is a graph depicting the *in vitro* effect of increasing concentrations of SL-  
11103 (●), SL-11104 (○), SL-11108 (■), and BE-444 (◆) on the survival of cultured  
25 human brain tumor cells U251 MG NCI.

ED<sub>50</sub> of SL-11103 = 9.5  $\mu$ M, SL-11104 = 14.71  $\mu$ M, SL-11108 = 2.0  $\mu$ M,  
and BE-444 = 0.2  $\mu$ M.

Fig. 23 is a graph depicting the *in vitro* effect of increasing concentrations of  
30

SL-11091 (●), SL-11094 (○), and BE-343 (■) on the survival of cultured human prostate cancer cells PC3.

ED<sub>50</sub> of SL-11091 >31.25 μM, SL-11094 >31.25 μM, and BE-343 = 0.24 μM.

Fig. 24 is a graph depicting the *in vitro* effect of increasing concentrations of SL-11091 (●), SL-11094 (○), and BE-343 (■) on the survival of cultured human prostate cancer cells DU145.

ED<sub>50</sub> of SL-11091 = 4.33 μM, SL-11094 = 15.4 μM, and BE-343 = 0.044 μM.

Fig.25 is a graph depicting the *in vitro* effect of increasing concentrations of SL-11091 (●), SL-11094 (○), and BE-343 (■) on the survival of cultured human colon cancer cells HT29.

ED<sub>50</sub> of SL-11091 >31.25 μM, SL-11094 = 28.8 μM, and BE-343 = 0.6 μM.

Fig. 26 is a graph depicting the *in vitro* effect of increasing concentrations of SL-11091 (●), SL-11094 (○), and BE-343 (■) on the survival of cultured human lung cancer cells A549.

ED<sub>50</sub> of SL-11091 >31.25 μM, SL-11094 >31.25 μM, and BE-343 = 0.2 μM.

Fig. 27 is a graph depicting the *in vitro* effect of increasing concentrations of SL-11091 (●), SL-11094 (○), and BE-343 (◆) on the survival of cultured human breast cancer cells MCF7.

ED<sub>50</sub> of SL-11091 >31.25 μM, SL-11094 >31.25 μM, and BE-343 = 0.5 μM.

Fig. 28 is a graph depicting the *in vitro* effect of increasing concentrations of SL-11091 (●), SL-11094 (○), and BE-343 (■) on the survival of cultured human brain tumor cells U251 MG NCI.

ED<sub>50</sub> of SL-11091 >31.25 μM, SL-11094 >31.25 μM, and BE-343 = 0.14 μM.

Fig. 29 is a graph depicting the *in vitro* effect of increasing concentrations of SL-11141(●), SL-11144 (□), SL-11150 (■) on the survival of cultured human prostate cancer cells PC3.

ED<sub>50</sub> of SL-11141>31.25 μM, SL-11144 = 0.3 μM, and SL-11150 = 0.5 μM.

Fig. 30 is a graph depicting the *in vitro* effect of increasing concentrations of SL-11141 (●), SL-11144 (□), SL-11150 (■) on the survival of cultured human prostate cancer cells DU145.

ED<sub>50</sub> of SL-11141 = 0.13 μM, SL-11144 = 0.1 μM, and SL-11150 = 0.11 μM.

**Fig. 31** is a graph depicting the *in vitro* effect of increasing concentrations of SL-11141 (●), SL-11144 (□), SL-11150 (■) on the survival of cultured human prostate cancer cells DUPRO.

ED<sub>50</sub> of SL-11141 = 0.71 μM, SL-11144 = 0.36 μM, and SL-11150 = 0.48 μM.

5        **Fig. 32** is a graph depicting the *in vitro* effect of increasing concentrations of SL-11141 (●), SL-11144 (□), SL-11150 (■) on the survival of cultured human prostate cancer cells LNCAP.

ED<sub>50</sub> of SL-11141 = 0.07 μM, SL-11144 = 0.20 μM, and SL-11150 = 0.23 μM.

10        **Fig. 33** illustrates synthetic methodology used to prepare the compounds of the invention.

**Fig. 34** illustrates additional synthetic methodology used to prepare the compounds of the invention.

**Fig. 35** illustrates additional synthetic methodology used to prepare the compounds of the invention.

15        **Fig. 36** illustrates additional synthetic methodology used to prepare the compounds of the invention.

**Fig. 37** illustrates additional synthetic methodology used to prepare the compounds of the invention.

20        **Fig. 38** illustrates additional synthetic methodology used to prepare the compounds of the invention.

**Fig. 39** illustrates additional synthetic methodology used to prepare the compounds of the invention.

**Fig. 40A** illustrates additional synthetic methodology used to prepare the compounds of the invention.

25        **Fig. 40B** illustrates additional synthetic methodology used to prepare the compounds of the invention.

**Fig. 41** illustrates synthetic methodology used to prepare porphyrin-polyamine conjugates of the invention.

30        **Fig. 42** illustrates additional synthetic methodology used to prepare porphyrin-polyamine conjugates of the invention.

**Fig. 43** illustrates additional synthetic methodology used to prepare porphyrin-polyamine conjugates of the invention.

Fig. 44 illustrates additional synthetic methodology used to prepare porphyrin-polyamine conjugates of the invention.

Fig. 45 is a graph depicting the *in vitro* effect of increasing concentrations of SL-11162 on the survival of human prostate cancer cells DuPro. The concentrations of SL-11162 are plotted in the X-axis and the corresponding surviving fractions of cells are plotted in the Y-axis.

Fig. 46 is a graph depicting the *in vitro* effect of increasing concentrations of SL-11184 on the survival of human prostate cancer cells DuPro. The concentrations of SL-11184 are plotted in the X-axis and the corresponding surviving fractions of cells are plotted in the Y-axis.

Fig. 47 is a graph depicting the *in vitro* effect of increasing concentrations of SL-11202 on the survival of human prostate cancer cells DuPro. The concentrations of SL-11202 are plotted in the X-axis and the corresponding surviving fractions of cells are plotted in the Y-axis.

Fig. 48 is a graph depicting the *in vitro* effect of increasing concentrations of SL-11184 on the survival of human prostate cancer cells PC-3. The concentrations of SL-11184 are plotted in the X-axis and the corresponding surviving fractions of cells are plotted in the Y-axis.

Fig. 49 is a graph depicting the *in vitro* effect of increasing concentrations of SL-11202 on the survival of human prostate cancer cells PC-3. The concentrations of SL-11202 are plotted in the X-axis and the corresponding surviving fractions of cells are plotted in the Y-axis.

## MODES FOR CARRYING OUT THE INVENTION

The present invention encompasses novel conformationally restricted polyamine analogs and compositions comprising these compounds. These analogs are useful as anti-proliferatives for suppressing cell proliferation. The analogs find use in treatment of various diseases, including use as anti-cancer agents in the treatment of various cancers. The analogs are also useful as anti-microbial agents. Novel polyamine analogs include the compounds shown in the synthetic schemes (Figures 33-40) and Table 1.

The invention also includes polyamine-porphyrin conjugates. These conjugates include the compounds depicted in Table 2 and those shown in the synthetic schemes (Figures 41-44).

## 5 Definitions

By "polyamine analog" is meant an organic cation structurally similar but non-identical to polyamines such as spermine and/or spermidine and their precursor, diamine putrescine. By a "polyamine", a term well-understood in the art, is meant any of a group of aliphatic, straight-chain amines derived biosynthetically from amino acids; polyamines are reviewed in Marton et al. (1995) *Ann. Rev. Pharm. Toxicol.* 35:55-91. Polyamine analogs can be branched or un-branched. Polyamine analogs include, but are not limited to, BE-4444 [1,19-*bis* (ethylamino)-5,10,15-triazanonadecane]; BE-333 [N1,N11-diethylnorspermine; DENSPM; 1,11-*bis* (ethylamino)-4,8-diazaundecane; thermine; Warner-Parke-Davis]; BE-33 [N1,N7-*bis*(ethyl) norspermidine]; BE-34 [N1,N8-*bis*(ethyl) spermidine]; BE-44 [N1,N9-*bis*(ethyl) homospermidine]; BE-343 [N1,N12-*bis*(ethyl) spermine; diethylspermine-N1-N12; DESPM]; BE-373 [N,N'-*bis* (3-ethylamino) propyl)-1,7-heptane diamine, Merrell-Dow]; BE-444 [N1,N14-*bis*(ethyl) homospermine; diethylhomospermine-N1-N14]; BE-3443 [1,17-*bis*(ethylamino)-4,9,14-triazaheptadecane]; and BE-4334 [1,17-*bis*(ethylamino)-5,9,13-triazaheptadecane]; 1,12-Me<sub>2</sub>-SPM [1,12-dimethylspermine]. Additional polyamine analogs are disclosed in International Patent Application WO 98/17624 and U.S. Patent No. 5,889,061. Various novel polyamine analogs are illustrated in the synthetic schemes in Figs. 33-40, and Table 1.

By "conformationally restricted" is meant that, in a polyamine analog, at least two amino groups in the molecule are locked or limited in spatial configuration relative to each other. The amino groups within the molecule may be primary, secondary, tertiary, or quaternary, and are preferably primary or secondary amino groups. The relative movement of two amino groups can be restricted, for example, by incorporation of a cyclic or unsaturated moiety between them (exemplified, but not limited to, a ring, such as a three-carbon ring, four-carbon ring, five-carbon-ring, six-carbon ring, or a double or triple bond, such as a double or triple carbon bond). Groups restricting conformational flexibility by means of steric hindrance, yet structurally favorable to the anti-proliferative, anti-cancer, or anti-microbial effects, can also be used according to the invention. A "conformationally



restricted” polyamine analog can comprise at least two amino groups which are conformationally restricted relative to each other, but can also further comprise amino groups which are not conformationally restricted relative to each other. Flexible molecules such as spermine and BE-444 can have a myriad of conformations and are therefore not conformationally restricted. Conformationally restricted polyamine analogs include, but are not limited to, the compounds listed in Table 1 and illustrated in Figures 33-40.

An “indication” includes any symptom or the like which points out a suitable remedy or treatment or which shows the presence of a disease or other unhealthful condition. As used herein, an “indication” also includes a “disease” itself, where a disease is a condition of an organ, part, structure or system of the body in which there is incorrect function resulting from the effect(s) of heredity, infection, diet and/or environment, and/or other causes. The indication can include cancer. By “cancer” is meant the abnormal presence of cells which exhibit relatively autonomous growth, so that they exhibit an aberrant growth phenotype characterized by a significant loss of cell proliferation control. Cancerous cells can be benign or malignant. In various embodiments, the cancer affects cells of the bladder, blood, brain, breast, colon, digestive tract, lung, ovaries, pancreas, prostate gland, or skin. In other embodiments, the indication can also include, but is not limited to, Alzheimer’s disease, epilepsy, multiple sclerosis, problems associated with tissue grafts and organ transplants, psoriasis, restenosis, stomach ulcers, or tissue overgrowth after surgery. In other embodiments, the indication is an infection or infestation of parasites, bacteria, fungi or insects.

An “individual” is a vertebrate, preferably a mammal, more preferably a human. Mammals include, but are not limited to, farm animals, sport animals, rodents, primates, and pets. Preferably, the individual is known or suspected to be afflicted by cancer or another disease treatable with a polyamine analog.

An “effective amount” or “therapeutic amount” is an amount sufficient to effect beneficial or desired clinical results. An effective amount can be administered in one or more administrations. For purposes of this invention, an effective amount of a polyamine analog is an amount that is sufficient to palliate, ameliorate, stabilize, reverse, slow or delay the progression of the disease state. A therapeutic amount of a polyamine of the present invention is an amount sufficient to inhibit proliferation of diseased cells. A polyamine analog is considered to be an effective anti-tumor or anti-cancer agent if it is effective

against at least one type of cancer cell line, even if it is not effective against a different cell line.

As used herein, "treatment" is an approach for obtaining beneficial or desired clinical results. For purposes of this invention, beneficial or desired clinical results include, but are not limited to, alleviation of symptoms, diminishment of extent of disease, stabilization (i.e., not worsening) of state of disease, prevention of spread (i.e., metastasis) of disease, delay or slowing of disease progression, amelioration or palliation of the disease state, improvement in quality of enjoyment of life, and remission (whether partial or total), whether detectable or undetectable. "Treatment" can also mean prolonging survival as compared to expected survival if not receiving treatment.

"Palliating" a disease means that the extent and/or undesirable clinical manifestations of a disease state are lessened and/or time course of the progression is slowed or lengthened, as compared to not administering polyamine analogs of the present invention. Preferred polyamine analogs for methods of treating and palliating a disease include the compounds illustrated in Table 1.

#### *Polyamine analogs of the present invention*

Novel polyamine analogs can be first designed based on current knowledge of polyamines and their analogs. Without wishing to be bound by any particular theory explaining polyamine analog toxicity, the inventors suggest that relevant knowledge includes that related to polyamine interaction with DNA and ability to induce structural changes in nucleic acids. Feuerstein et al. (1991); Gosule et al. (1978) *J. Mol. Biol.* 121:311-326; Behe et al. (1981) *Proc. Natl. Acad. Sci. USA* 78:1619-23; Jain et al. (1989) *Biochem.* 28:2360-2364; and Basu et al. (1990) *Biochem. J.* 269:329-334. Alternatively, a novel polyamine analog can be designed based on its likely ability to inhibit cell growth by suppressing natural polyamine synthesis or depleting the intracellular natural polyamine pool. Porter et al. (1988) in *Advances in Enzyme Regulation*, Pergamon Press, pp. 57-79. Preferably, the novel polyamine analog is conformationally restricted. In the next step, the polyamine analog is tested *in vitro* for efficacy in inhibiting proliferation of diseased cells, such as tumor cells. If the polyamine analog passes this test, it can then be tested in animals, such as nude mice with cancer xenografts. If the compound is found to be efficacious, testing can then proceed to human trials.

The present invention encompasses novel polyamine analogs, such as those shown in Figs. 33-40 and Table 1. Polyamine analogs of the present invention are conformationally restricted. Conformation is a determinant of the spatial arrangement of the pharmacophore or functional groups which interact with receptor binding sites. The latter prefer specific ligand conformations or a specific distribution of conformations. A flexible molecule such as spermine or BE-4444 can have a myriad of conformations. The conformer that binds to the macromolecule (e.g., DNA or RNA) may not necessarily be the one with the lowest energy as determined by spectroscopic methods or theoretically by molecular mechanics calculations. The binding energy of the polyamine binding to the nucleic acid may be overcome with formation of an unstable conformer. Conversely, in the presence of a conformationally rigid analog of a flexible molecule, the host macromolecule might change its overall conformation or the distances from one strand to the other. Hydrogen bonding is the main binding force of either spermine or spermidine associating with the helical region of a tRNA, and very likely also with DNA. Frydman et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:9186-9191; and Fernandez et al. (1994) *Cell Mol. Biol.* 40: 933-944. The secondary amino groups present in the linear spermine analogs BE-343, BE-444, and BE-4444 are the groups most directly involved in the formation of the hydrogen bonds with the paired bases of tRNA. Therefore, these amino groups that usually flank the central four-carbon or three-carbon segment of the polyamine analog can be considered as the pharmacophore of the polyamine analog. When the nitrogens are separated by only a two-carbon segment they are not protonated at pH 7.4 and hence they do not form hydrogen bonds. If these amino groups are locked into various configurations by the incorporation of cyclic or unsaturated moieties into the polyamine analog molecule, a conformationally rigid analog is obtained. When such analogs bind to DNA or tRNA, they will very likely induce a change in the conformation of the nucleic acid strands or loops that may differ from the conformational changes induced by the natural polyamines. A series of conformationally restricted analogs of *bis*-ethyl-spermine that exhibited cytotoxic activities are illustrated in Table 1 and described below.

#### *Porphyrin-polyamine conjugates of the present invention*

The present invention also encompasses novel porphyrin-polyamine conjugates such as those depicted in Figs. 41-44 and in Table 3. These conjugates combine the

selective uptake of porphyrins by tumors with the cytotoxic and cytostatic effects of polyamine analogs. By "porphyrin-polyamine conjugate" is meant any porphyrin compound and any polyamine compound, linked by at least one, and preferably only one, covalent bond. Porphyrin-polyamine conjugates can incorporate any polyamine, whether that polyamine is conformationally restricted or not. Preferably, the polyamine conjugate is conformationally restricted. By "porphyrin" is meant any compound incorporating the porphine ring or derivatives of the porphine ring. Preferred covalent linkages include amide linkages and amine linkages. Examples of porphyrin-polyamine conjugates are depicted in Table 2.

#### *Polyamine analogs and inhibition of cell growth*

Polyamine analogs of the present invention are likely to be useful for treatment of a variety of diseases, including cancer, Alzheimer's disease, epilepsy, multiple sclerosis, problems associated with tissue grafts and organ transplants, psoriasis, restenosis, stomach ulcers, or tissue overgrowth after surgery, or an infection or infestation of parasites, bacteria, fungi or insects. In order to evaluate the efficacy of a particular novel polyamine for a particular medicinal application, the compounds can be first tested against appropriately chosen test cells *in vitro*. In a non-limiting example, polyamine analogs can be tested against tumor cells, for example, prostate tumor cells. On the basis of the unique nature of polyamine metabolism in the prostate gland, exemplary experiments can utilize cell lines capable of growing in culture as well as *in vivo* in athymic nude mice, such as LNCaP. Horoszewicz et al. (1983) *Cancer Res.* 43:1809-1818. Culturing and treatment of carcinoma cell lines, cell cycle and apoptosis determinations based on flow cytometry; enzyme assays including ODC, SAMDC and SSAT activities; and high pressure liquid chromatography detection and quantitation of natural polyamines and polyamine analogs are described in the art, for example, Mi et al. (1998) *Prostate* 34:51-60; Kramer et al. (1997) *Cancer Res.* 57:5521-27; and Kramer et al. (1995) *J. Biol. Chem.* 270:2124-2132. Evaluations can also be made of the effects of the novel polyamine analog on cell growth and polyamine-related metabolism. Analysis begins with IC<sub>50</sub> determinations based on dose-response curves ranging from 0.1 to 1000  $\mu$ M performed at 72 hr. From these studies, conditions can be defined which produce about 50% growth inhibition and used to: (a) follow time-dependence of growth inhibition for up to 6 days, with particular attention to

decreases in cell number, which may indicate drug-induced cell death; (b) characterize analog effects on cell cycle progression and apoptosis using flow cytometry (analysis to be performed on attached and detached cells); (c) examine analog effects on polyamine metabolic parameters, including the biosynthetic enzymes ODC, SAMDC, the catabolic enzyme SSAT and polyamine pools themselves. Analog effects can be normalized to intracellular concentrations (by HPLC analysis), which also provide an indication of their relative ability to penetrate cells. Marked differences in analog uptake can be further characterized by studying analog ability to utilize and regulate the polyamine transporter, as assessed by competition studies using radiolabeled spermidine, as previously described in Mi et al. (1998).

*In vivo testing of polyamine analogs and porphyrin-polyamine conjugates*

Analog and conjugates found to have potent or mechanism-based anti-proliferative activity *in vitro* towards cultured carcinoma cells can be evaluated in *in vivo* model systems. The first goal is to determine the relative toxicity of the analogs or conjugates in non-tumor-bearing animals, such as DBA/2 mice. Groups of three animals each can be injected intraperitoneally with increasing concentrations of a compound, beginning at, for example, 10 mg/kg. Toxicity as indicated by morbidity is closely monitored over the first 24 hr. A well-characterized polyamine analog, such as BE-333, can be used as an internal standard in these studies, since a data base has already been established regarding acute toxicity via a single dose treatment relative to chronic toxicity via a daily x 5 d schedule. Thus, in the case of new analogs, single dose toxicity relative to BE-333 is used to project the range of doses to be used on a daily x 5 d schedule.

After the highest tolerated dosage on a daily x 5 d schedule is deduced, antitumor activity is determined. Typically, tumors can be subcutaneously implanted into nude athymic mice by trocar and allowed to reach 100-200 mm<sup>3</sup> before initiating treatment by intraperitoneal injection daily x 5 d. Most analogs or conjugates can be given in a range between 10 and 200 mg/kg. Analog and conjugates can be evaluated at three treatment dosages with 10-15 animals per group (a minimum of three from each can be used for pharmacodynamic studies, described below). Mice can be monitored and weighed twice weekly to determine tumor size and toxicity. Tumor size is determined by multi-directional measurement from which volume in mm<sup>3</sup> is calculated. Tumors can be followed until

median tumor volume of each group reaches 1500 mm<sup>3</sup> (i.e., 20% of body weight), at which time the animals can be sacrificed. Although the initial anti-tumor studies focuses on a daily x 5 d schedule, constant infusion can be performed via Alzet pump delivery for 5 days since this schedule dramatically improves the anti-tumor activity of BE-333 against A549 human large cell lung carcinoma. Sharma et al. (1997) *Clin. Cancer Res.* 3:1239-1244. In addition to assessing anti-tumor activity, free analog levels or conjugate levels in tumor and normal tissues can be determined in test animals.

*Methods of administration of polyamine analogs and porphyrin-polyamine conjugates*

The polyamine analogs or porphyrin-polyamine conjugates of the present invention can be administered to an individual via any route known in the art, including, but not limited to, those disclosed herein. Preferably administration of the novel polyamine analogs or porphyrin-polyamine conjugates is by intravenous routes. Other methods of administration include but are not limited to, oral, intrarterial, intratumoral, intramuscular, subcutaneous, intraperitoneal, gastrointestinal, and directly to a specific or affected organ. The novel polyamine analogs and porphyrin-polyamine conjugates described herein are administratable in the form of tablets, pills, powder mixtures, capsules, injectables, solutions, suppositories, emulsions, dispersions, food premixes, and in other suitable forms. Additional methods of administration are known in the art. The pharmaceutical dosage form which contains the compounds described herein is conveniently admixed with a non-toxic pharmaceutical organic carrier or a non-toxic pharmaceutical inorganic carrier. Typical pharmaceutically-acceptable carriers include, for example, mannitol, urea, dextrans, lactose, potato and maize starches, magnesium stearate, talc, vegetable oils, polyalkylene glycols, ethyl cellulose, poly(vinylpyrrolidone), calcium carbonate, ethyl oleate, isopropyl myristate, benzyl benzoate, sodium carbonate, gelatin, potassium carbonate, silicic acid, and other conventionally employed acceptable carriers. The pharmaceutical dosage form can also contain non-toxic auxiliary substances such as emulsifying, preserving, or wetting agents, and the like. A suitable carrier is one which does not cause an intolerable side effect, but which allows the novel polyamine analogs or porphyrin-polyamine conjugates to retain its pharmacological activity in the body. Formulations for parenteral and nonparenteral drug delivery are known in the art and are set forth in *Remington's Pharmaceutical Sciences*, 18th Edition, Mack Publishing (1990).

Solid forms, such as tablets, capsules and powders, can be fabricated using conventional tableting and capsule-filling machinery, which is well known in the art. Solid dosage forms can contain any number of additional non-active ingredients known to the art, including excipients, lubricants, dessicants, binders, colorants, disintegrating agents, dry flow  
5 modifiers, preservatives, and the like. Liquid forms for ingestion can be formulated using known liquid carriers, including aqueous and non-aqueous carriers, suspensions, oil-in-water and/or water-in-oil emulsions, and the like. Liquid formulations can also contain any number of additional non-active ingredients, including colorants, fragrance, flavorings, viscosity modifiers, preservatives, stabilizers, and the like. For parenteral administration,  
10 novel polyamine analogs or porphyrin-polyamine conjugates can be administered as injectable dosages of a solution or suspension of the compound in a physiologically acceptable diluent or sterile liquid carrier such as water or oil, with or without additional surfactants or adjuvants. An illustrative list of carrier oils would include animal and vegetable oils (peanut oil, soy bean oil), petroleum-derived oils (mineral oil), and synthetic  
15 oils. In general, for injectable unit doses, water, saline, aqueous dextrose and related sugar solutions, and ethanol and glycol solutions such as propylene glycol or polyethylene glycol are preferred liquid carriers. The pharmaceutical unit dosage chosen is preferably fabricated and administered to provide a final concentration of drug at the point of contact with the cancer cell of from 1  $\mu$ M to 10 mM. More preferred is a concentration of from 1  
20 to 100  $\mu$ M. As with all pharmaceuticals, the optimal effective concentration of novel polyamine analogs or porphyrin-polyamine conjugates will need to be determined empirically and will depend on the type and severity of the disease, route of administration, disease progression and health and mass or body area of the patient. Such determinations are within the skill of one in the art. Polyamine analogs or porphyrin-polyamine conjugates  
25 can be administered as the sole active ingredient, or can be administered together, or in combination with another active ingredient, including, but not limited to, cytotoxic agents, antibiotics, antimetabolites, nitrosourea, vinca alkaloids, polypeptides, antibodies, cytokines, etc.

30 The following examples are provided to illustrate but not limit the invention.

## EXAMPLES

## Synthesis of conformationally-restricted polyamine analogs and of porphyrin-polyamine conjugates

5                   a)       **Spermine and homospermine analogs containing a conformational restriction.**

                  Scheme 2 exemplifies a N<sup>α</sup>, N<sup>ω</sup>-bisethyl homospermine analog **7** containing a central *trans*-unsaturated bond. Amide **4** was prepared as described in Scheme 1 by alkylation of amide **1** with bromobutyronitrile to give **2**, followed by reduction of the nitrile  
10       to the amine **3** that was mesitylsulfonated to **4**. *Trans*-allylic diester **5** was used to alkylate amide **4** and the tetramide **6** was obtained. Deprotection gave the *trans*-tetramide **7** (Scheme 2).

                  Introduction of a triple bond in the butane segment of homospermine also reduces its mobility. This was achieved by starting with the butyne diester **8** and following the  
15       sequence of reactions outlined above (Scheme 3). Schemes 15-20 are further examples of the synthesis of polyamine spermine and homospermine analogs of this type.

b)       **Synthesis of pentamines with conformational restrictions.**

                  Schemes 4-14 are outlines of the syntheses of conformationally restricted pentamines. Scheme 4 depicts the reaction of *cis*-1-chloro-4-phthalimido butene with  
20       amide **1** to give **11**. Hydrazinolysis of **11** gave **12** which was amidated to **13**. Reaction of the latter with 1,4-diiodobutane gave **14**, while reaction with equimolar amounts of *cis*-1,4-dichlorobutene gave **15**.

                  Amide **4** was alkylated with either 4-chlorobutyronitrile to give **16** or with *cis*-1,4-dichlorobutene to give **19**. Nitrile **16** was reduced with hydrogen over Ni Raney to  
25       the amine **17** and the latter transformed in to the amide **18** (Scheme 5). Condensation of **18** with the chloroalkyl intermediate **15** gave the pentamide **20** that was deprotected to the pentamine **21** (Scheme 6). Condensation of **18** with the iodoalkyl derivative **14** gave **22** that was deprotected to the pentamine **23** (Scheme 7). Condensation of **18** and **19** gave pentamide **24** that was deprotected to the pentamine **25** (Scheme 8). Using **14** as the  
30       alkylating agent, mesitylenesulfonamide was dialkylated to give **26**, and the latter deprotected to give **27** (Scheme 9). The analogous reaction carried out using **15** as alkylating agent, gave **28** and after deprotection led to the pentamine **29** (Scheme 10).



Alkylation of mesitylenesulfonamide with **19** gave the pentamide **30**, which was deprotected to **31** (Scheme 11). When **19** was used to alkylate an equimolar amount of mesitylenesulfonamide then **32** was obtained. Alkylation of **32** with **14** gave **33**, that was deprotected to give **34** (Scheme 12). When the chloroalkyl intermediate **15** was used to alkylate one equivalent of mesitylenesulfonamide, then the triamide **35** was obtained. Reaction of **35** with **14** gave **36** which was then deprotected to **37** (Scheme 13). Condensation of **35** and **19** gave the pentamide **38** that was deprotected to **39** (Scheme 14). The above mentioned Schemes describe the synthesis of *cis*-compounds. The same synthetic methodology can be used to obtain the *trans*-isomers, or *cis* and *trans* bonds in different segments within the same molecule.

**c) Polyamine analogs with diamidine substituents.**

A new class of polyamine analogs is shown in Scheme 21. They derive from 1,4-dibenzylputrescine, 1,5-dibenzylcadaverine, and 1,6-dibenzylhexanediamine. They are diamidine derivatives, where the diamidine residues are carrier groups that have been shown to be efficient in the transport of drugs into different protozoa. The general procedure of synthesis was based on the condensation of 4-cyanobenzaldehyde with the diaminoalkanes to give the Schiff bases, followed by reduction *in situ* to the corresponding dinitriles **68**. The latter were converted to the diamidines **69** through their iminoethers.

**d) Synthesis of oligoamines.**

Scheme 22 describes the synthesis of a N-2 hydroxyethyl derivative of a pentamine such as **75**. Starting with **18**, alkylation with 4-bromobutyronitrile gave **70**. Reduction of the nitrile of **70** and mesitylenesulfonylation of the resulting amino group gave **71**. It was alkylated again with 4-bromobutyronitrile to give **72**, and again reduced and mesitylsulfonylated to give **73**. The latter was then alkylated with the benzyl ester of 2-bromoethanol to give **74**. Treatment with hydrobromic acid in acetic acid cleaved both the mesitylene sulfonyl protecting groups and the benzyl ether residue to give **75**.

Scheme 23 reports the synthesis of a *trans*-decamine **77** and of a *cis*-decamine **79**. Starting with the pentamide **73** (Scheme 22) and by reaction with *trans*-diester **5** (Scheme 2) the decamide **76** was prepared, which on deprotection gave **77** as a decahydrochloride. In an analogous manner, by condensation of **73** with the *cis*-1,4-

dimesityleneoxy-2-butene, the decamide **78** was prepared, which on deprotection gave **79** as a decahydrochloride.

Scheme 24 outlines the synthesis of a N-2 hydroxyethyl *trans*-decamine **92** and a *cis*-2-hydroxyethyl decamine **95**. The procedure repeats almost all the procedures described in the foregoing schemes. The synthesis of **80** proceeded by alkylating BOC-mesitylenesulfonamide with the benzyl ester of 2-bromoethanol. Cleavage of the BOC protecting group leads to **81**, alkylation with 4-bromobutyronitrile then gave **82**, and after reduction of the nitrile group and reaction with mesitylene sulfonyl chloride the diamide **83** was obtained. Again, alkylation with 4-bromobutyronitrile led to **84**, reduction and mesitylsulfonylation gave **85**, alkylation of **85** gave **86**, reduction and mesitylsulfonylation gave **87**, and alkylation, reduction and mesitylsulfonylation performed on **87** gave **89**. Alkylation of **73** with *trans*-1,4-dibromo-2-butene gave **90**. Alkylation of **89** with **90** gave **91**, which after deprotection gave the *trans*- $\omega$ -hydroxy-decamine **92**. Alkylation of **73** with *cis*-1,4-dichloro-2-butene gave **93**. Alkylation of **89** with **93** gave **94**. Deprotection of **94** gave the *cis*- $\omega$ -hydroxy-decamine **95**, isomeric with **92**.

**e) Synthesis of porphyrin-polyamine conjugates.**

Scheme 25 outlines the synthesis of porphyrin-polyamine. Starting with the above mentioned amide **18** (Scheme 5), alkylation with the benzyl ether of 4-bromobutanol gave **96**. Treatment with acid cleaved both the sulfonate and the benzyl protecting groups to give **97**. Protection of the free amino residues with Boc followed by Mitsunobu reaction using phthalimide gave **99**. Cleavage of the phthalyl protecting group with hydrazine hydrate gave polyamine **100**. It was condensed with mesoporphyrin IX to give **101**, and after cleavage of the Boc protecting group **102** (SL-11162) was obtained.

Scheme 26 outlines the synthesis of a porphyrin-bisethylpolyamine conjugate. The synthesis of the intermediate **105** follows the pattern described above. The primary alcohol is oxidized to the aldehyde **106** in Swern oxidation reaction. The aldehyde **106** was then condensed with ethylamine using a reductive alkylation reaction that gave **107**. The latter was coupled with mesoporphyrin IX using amide forming procedures and the resulting **108** was deprotected in acid to **109** (SL-11184).

Scheme 27 illustrates on the preparation of a porphyrin-polyamine conjugate, where an amine group tethers the polyamine to the porphyrin. The amide derivative **111** was

prepared as described above and was then reduced using metal hydrides and deprotected to yield **112**. The synthesis of the porphyrin-polyamine **112** could also be achieved as shown in the Scheme 28. Reduction of the diester of mesoporphyrin IX under controlled conditions gave the dialdehyde **113**. Reductive alkylation of **100** with **113** followed by deprotection also gave **112** (SL-11202).

## EXAMPLE 1

### Synthesis Of Polyamine Compounds

**Compound 2:** NaH (80%, 1.08 g, 36 mmol) was added to a solution of amide **1** (6.81 g, 30 mmol) in DMF (50 ml) in an ice-water bath under N<sub>2</sub>. The mixture was stirred for 1 h and a solution of 4-bromobutyronitrile (4.88 g, 33 mmol) in DMF (10 ml) was added in portions. The mixture was stirred over night at 75°C. The solvent was distilled off, the residue taken up in chloroform washed with a saturated solution of ammonium chloride, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The residue was purified by flash chromatography on silica gel (hexane/ethyl acetate 3:1) to yield 8.0 g (90%) of **2** as a colorless oil. <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 1.05 (t, 3H), 1.90 (m, 2H), 2.30 (b, m, 5H), 2.60 (s, 6H), 3.20 (q, 2H), 3.35 (t, 2H), 6.95 (s, 2H); <sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 12.50, 20.61, 22.43, 23.60, 31.05, 36.12, 40.39, 43.78, 118.62, 131.79, 132.67, 139.71, 142.41. MS-EI (m/z) 294 (M<sup>+</sup>).

**Compound 4:** Nitrile **2** (7.8 g, 27 mmol) was dissolved in a mixture of ethanol (150 ml) and concentrated hydrochloric acid (1.5 ml). PtO<sub>2</sub> was added (700 mg) and the mixture was hydrogenated at 50 psi over night. The catalyst was filtered off and the solvent evaporated. The residue (78 g, 98%) was used in the next step without further purification. The free base gave <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 1.00 (t, 3H), 1.55 (m, 4H), 2.25 (s, 3H), 2.80 (t, 2H), 3.20 (m, 4H), 6.95 (s, 2H); <sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 12.54, 20.69, 22.53, 24.72, 27.65, 39.92, 40.29, 44.59, 131.71, 133.21, 139.82, 142.09. FAB-MS (m/z) 299 (M<sup>+</sup> + 1). Mesitylenesulfonyl chloride (8.8 g, 40.5 mmol) in dioxane (30 ml) was added dropwise to a stirred mixture of compound **3** (7.8 g, 27 mmol) dissolved in dioxane (60 ml) and 50% KOH (30 ml) at 0°C. The reaction mixture was allowed to reach 20°C and then kept over night. An excess of water was added and the mixture was extracted with chloroform, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The oily residue was crystallized from ethyl acetate/hexane yielding **4**; 10.9 g (82%); mp 71.5-72°C. <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 1.00 (t, 3H), 1.10-1.50 (m,

4H), 2.30 (s, 6H), 2.55, 2.60 (s, 12H), 2.85 (q, 2H), 3.15 (m, 4H), 4.70 (t, 1H), 6.95, 7.00 (s, 4H);  $^{13}\text{C}$ -NMR ( $\text{CDCl}_3$ ):  $\delta$  12.70, 20.92, 21.04, 22.73, 22.92, 24.58, 26.68, 40.04, 42.02, 44.42, 131.91, 133.31, 133.64, 138.99, 140.05, 142.15, 142.35. MS-FAB ( $m/z$ ) 480 ( $\text{M}^+$ ).

5 **Compound 5: (E)-2-Butene-1,4-diyl bis[mesitylenesulfonate]** : (E)-2-Butene-1,4-diol (1.76 g, 20 mmol), and benzyltriethylammonium bromide (270 mg, 1 mmol) were dissolved in 30 ml of a 50% potassium hydroxide solution and 30 ml of dioxane. The mixture was stirred at  $5^\circ\text{C}$  and mesitylenesulfonyl chloride (8.72 g, 40 mmol) dissolved in 30 ml of dioxane was added dropwise. When the addition was over, stirring was continued  
10 for 1 h, water was then added, and the white precipitate was filtered and crystallized from chloroform-hexane to yield **5**; 7.0 g (77%); mp  $119\text{--}120^\circ\text{C}$ .  $^1\text{H}$ -NMR ( $\text{CDCl}_3$ ):  $\delta$  2.35 (s, 6H), 2.60 (s, 12H), 4.45 (d, 4H), 5.75 (b, 2H), 6.95 (s, 4H);  $^{13}\text{C}$ -NMR ( $\text{CDCl}_3$ ):  $\delta$  20.96, 22.52, 67.96, 127.67, 131.69, 131.74, 139.79, 143.45. MS-EI ( $m/z$ ), 452 ( $\text{M}^+$ ), 253, 200, 183. Anal. Calcd for  $\text{C}_{22}\text{H}_{28}\text{O}_6\text{S}_2$ : C, 58.40; H, 6.19. Found: C, 58.35; H, 6.22.

15 **Compound 6** was synthesized from **5** according to a procedure described elsewhere (Reddy et al., *J. Med. Chem.* 41:4723 (1998)) in 56% yield.  $^1\text{H}$ -NMR ( $\text{CDCl}_3$ ):  $\delta$  0.95 (t,  $J=7.15$  Hz, 6H,  $\text{CH}_3$ ), 1.34 (m, 8H,  $\text{CH}_2$ ), 2.29 (s, 12H,  $\text{CH}_3$ ), 2.55 (s, 24H,  $\text{CH}_3$ ), 3.09 (m, 12H,  $\text{NCH}_2$ ), 3.72 (d,  $J=4.53$  Hz, 4H,  $\text{NCH}_2$ ), 5.48 (t,  $J=4.31$  Hz, 2H,  $\text{CH}=\text{CH}$ ), 6.92 (s, 4H, Ph), 6.93 (s, 4H, Ph);  $^{13}\text{C}$ -NMR ( $\text{CDCl}_3$ ):  $\delta$  12.71, 20.90, 22.71, 22.76, 24.74, 40.04,  
20 42.21, 44.56, 45.69, 128.45, 131.88, 132.02, 140.05, 140.16, 142.20, 142.58. MS-FAB ( $m/z$ ) 1012.8 ( $\text{M}^+$ , 100%), 828.7, 646.7, 561, 176.

**Compound 7** was obtained from **6** as described elsewhere (Reddy et al., *J. Med. Chem.* 41:4723 (1998)) in 75% yield, mp  $>230^\circ\text{C}$ .  $^1\text{H}$ -NMR ( $\text{D}_2\text{O}$ ):  $\delta$  1.26 (t,  $J=12.5$  Hz, 6H,  $2\text{CH}_3$ ), 1.79 (m, 8H,  $\text{CH}_2$ ), 3.12 (m, 12H,  $\text{NCH}_2$ ), 3.80 (d,  $J=7.16$ , 4H,  $\text{NCH}_2$ ), 6.10 (m, 2H,  $\text{CH}=\text{CH}$ );  $^{13}\text{C}$ -NMR ( $\text{D}_2\text{O}$ ):  $\delta$  12.79, 25.10, 45.19, 48.53, 48.62, 50.36, 130.66. MS-MALDI ( $m/z$ ): 285.3 ( $\text{MH}^+$ , 100%).

**Compound 8** was obtained from the commercially available butyne diol.

Mesitylenesulfonyl chloride (19.5 g, 90 mmol) in dioxane (30 ml) was added dropwise to a stirred and cooled mixture of butyne diol (2.58 g, 30 mmol), 50% potassium hydroxide (30 ml) and triethylbenzyl ammonium bromide (405 mg, 1.5 mmol). Once the addition was  
30 over, the mixture was stirred at room temperature for an additional 3 h. An excess of water was added and the white precipitate was cooled over night, filtered off and dried.

Recrystallization from ethyl acetate/hexane afforded 8.6 g (63%) of **8**; mp 105-106°C. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.30 (s, 6H), 2.60 (s, 12H), 4.50 (s, 4H), 6.95 (s, 4H); <sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 20.93, 22.48, 56.13, 80.41, 130.65, 131.67, 139.98, 143.67. MS-EI (m/z) 450 (M<sup>+</sup>).

**Compound 9** was obtained following a procedure analogous to that described for compound **42** (see below). From 450 mg (1 mmol) of diester **8** and 1.05 g (2.2 mmol) of diamide **4**, 570 mg (56%) of tetramide **9** was obtained. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.90 (t, 6H), 1.30 (bs, 8H), 2.20 (s, 12H), 2.45 (s, 24H), 3.05 (m, 12H), 3.75 (s, 4H), 6.87 (s, 8H); <sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 12.70, 20.78, 22.68, 34.65, 39.97, 44.46, 44.99, 78.62, 131.85, 131.98, 132.34, 140.14, 142.13, 142.55. MS-FAB (m/z) 1010 (M<sup>⊕</sup>).

**Compound 10** was obtained following a procedure analogous to that described for compound **43** (see below). From 500 mg (0.49 mmol) of tetramide **9**, 160 mg (76%) of the tetrahydrochloride **25** was obtained; mp >280°C (decomp). <sup>1</sup>H-NMR (D<sub>2</sub>O): δ 1.30 (t, 6H), 1.80 (b, 8H), 2.90-3.25 (m, 12H), 4.05 (s, 4H); <sup>13</sup>C-NMR (D<sub>2</sub>O): δ 13.39, 25.64, 39.26, 45.72, 49.00, 49.20, 81.20. MS-MALDI 283 (M<sup>+</sup> + 1).

**Compound 11:** Mesitylenesulfonylethylamide **1** (3.1 g, 13.65 mmol) was dissolved in anhydrous DMF (30 ml) followed by the addition of NaH (85%, 0.423 g) in several portions. The mixture was stirred at room temperature for 1 h. N-(4-chloro-2-butenyl)-phthalimide (Aldrich, 3.06 g, 13 mmol) in 20 ml of DMF was added to the flask followed by stirring at 80°C over night. The mixture was cooled to room temperature, quenched with H<sub>2</sub>O (10 ml), and the solution was evaporated to dryness in vacuo. The solid residue was partitioned between 25 ml H<sub>2</sub>O and 25 CHCl<sub>3</sub>. The aqueous layer was extracted with CHCl<sub>3</sub> (3 x 25 ml), the organic layers were washed with brine (35 ml), dried (MgSO<sub>4</sub>), the solvent was evaporated to afford a gum which solidified upon trituration with hexane to give **11**. The <sup>1</sup>H-NMR and <sup>13</sup>C-NMR spectra showed that **11** was pure enough to be used in the next step without further purification, yield 4.75 g. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.16 (t, J=7.11 Hz, 3H, CH<sub>3</sub>), 2.29 (s, 3H, CH<sub>3</sub>), 2.63 (s, 6H, 2CH<sub>3</sub>), 3.29 (q, J=7.11 Hz, 2H, CH<sub>2</sub>), 4.06 (d, J=5.24 Hz, 2H, NCH<sub>2</sub>), 4.26 (d, J=5.72 Hz, 2H, NCH<sub>2</sub>), 5.59 (m, 2H, CH=CH), 6.95 (s, 2H, Ph), 7.71 (m, 2H, Ph), 7.83 (m, 2H, Ph); <sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 13.06, 20.89, 22.72, 34.35, 40.68, 42.01, 123.27, 126.69, 129.47, 131.90, 134.00, 140.24.

**Compound 12:** Amide 11 (20 g, 46.95 mmol) was dissolved in methanol, hydrazine monohydrate (5 ml, 98.52 mmol) was added and the solution stirred at 55°C for 24 h. Initially it was a homogeneous solution; however, after several hours a white solid precipitated. The mixture was cooled to room temperature, 300 ml of conc. HCl were added slowly (exothermic reaction), and stirring at room temperature was continued for 12 h more. Methanol was evaporated, and the resulting solid was extracted with CHCl<sub>3</sub> (3 x 150 ml). The aqueous layer was neutralized with 50% NaOH, extracted again with CHCl<sub>3</sub> (3 x 100 ml), the combined organic layers were dried (MgSO<sub>4</sub>); the solution was evaporated to afford a gum, which solidified under high vacuum to give **12**; yield 9.0 g (65%). The compound was purified by column chromatography using hexane, ethyl acetate (7:3) as eluent; mp 167-169°C. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.0 (t, J=7.1 Hz, 3H, CH<sub>3</sub>), 2.28 (s, 3H, CH<sub>3</sub>), 2.56 (s, 6H, 2CH<sub>3</sub>), 2.62 (br, NH<sub>2</sub>), 3.12 (q, J=7.1 Hz, 2H, NCH<sub>2</sub>), 3.73 (br, 2H, NCH<sub>2</sub>), 3.94 (d, J=6.0 Hz, 2H, NCH<sub>2</sub>), 5.80 (m, 2H, CH=CH), 6.92 (s, 2H, Ph); <sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 12.97, 20.93, 22.74, 36.43, 40.94, 42.08, 124.29, 131.89, 132.00, 132.62, 140.21, 142.67.

**Compound 13** was obtained from **12** as described for **4** in 96% yield. It was purified by column chromatography using hexane and ethyl acetate (4:1.5) as eluants; mp 98-99°C; <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.93 (t, J=5.85 Hz, 3H, CH<sub>3</sub>), 2.23 (s, 3H, CH<sub>3</sub>), 2.24 (s, 3H, CH<sub>3</sub>), 2.50 (s, 6H, 2CH<sub>3</sub>), 2.56 (s, 6H, 2CH<sub>3</sub>), 3.06 (q, J=7.15 Hz, 2H, NCH<sub>2</sub>), 3.48 (t, J=5.99 Hz, 2H, NCH<sub>2</sub>), 3.68 (d, J=5.72 Hz, 2H, NCH<sub>2</sub>), 4.58 (t, J=6.24 Hz, 1H, NH), 5.44 (m, 2H, CH=CH), 6.87 (s, 2H, Ph), 6.89 (s, 2H, Ph); <sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 12.80, 20.89, 22.64, 22.89, 39.01, 40.59, 41.41, 128.14, 128.46, 131.91, 131.96, 139.08, 140.19, 142.26, 142.54. MS-FAB (m/z) 479.2 (M<sup>+</sup>, 65%), 296.2, 279.1, 267.2, 183.1.

**Compound 15:** Amide 13 (4.79 g, 10 mmol) was dissolved in anhydrous DMF (40 ml) followed by addition of NaH (0.37 g) in several portions, the mixture stirred at room temperature for 2 h, cis-1,4-dichloro-2-butene (7.5 g, 60 mmol) in 10 ml DMF was added at once, and stirring was continued at 50°C over night. The mixture was cooled to room temperature, quenched with 10 ml H<sub>2</sub>O, the solvents were evaporated, and the contents were partitioned between H<sub>2</sub>O (50 ml) and CHCl<sub>3</sub> (50 ml). The aqueous layer was extracted with CHCl<sub>3</sub> (3 x 50 ml), the pooled organic layers were dried (MgSO<sub>4</sub>), evaporated, and **15** was purified by column chromatography using hexane, ethyl acetate (8.5:1.5) as eluants; yield 5.5 g (97%), mp 106-108°C. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.03 (t, J=7.33

Hz, 3H, CH<sub>3</sub>), 2.30 (s, 6H, 2CH<sub>3</sub>), 2.57 (s, 12H, 4CH<sub>3</sub>), 3.17 (q, J=7.31 Hz, NCH<sub>2</sub>), 3.71 (m, 4H, NCH<sub>2</sub>), 3.81 (d, J=6.87 Hz, 2H, NCH<sub>2</sub>), 3.95 (d, J=7.70 Hz, 2H, CHCl<sub>2</sub>), 5.50 (m, 3H, CH=CH), 5.74 (m, 1H, CH=CH), 6.93 (s, 2H, Ph), 6.95 (s, 2H, Ph); <sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 12.91, 22.70, 22.74, 38.20, 40.45, 41.60, 42.11, 42.33, 128.17, 128.95, 129.34, 129.40, 131.94, 132.08, 140.23, 140.34, 142.91. MS-FAB (m/z) 566.7 (M<sup>+</sup>, 100%), 153.4, 96.3.

**Compound 14** was prepared from **13** and 1,4-diiodobutane as described above for **15**. The product was purified by column chromatography using hexanes and ethyl acetate (4:1) as eluant; yield 79%. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.04 (t, J=7.10 Hz, 3H, CH<sub>3</sub>), 1.63 (m, 4H, CH<sub>2</sub>), 2.30 (s, 6H, 2CH<sub>3</sub>), 2.58 (s, 12H, 4CH<sub>3</sub>), 3.04 (t, J=6.50 Hz, 2H, CH<sub>2</sub>I), 3.16 (m, 4H, NCH<sub>2</sub>), 3.78 (d, J=5.14 Hz, 4H, NCH<sub>2</sub>), 5.55 (m, 2H, CH=CH), 6.94 (s, 2H, Ph), 6.95 (s, 2H, Ph); <sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 5.69, 12.92, 20.95, 22.72, 22.78, 28.25, 30.36, 40.47, 41.59, 42.11, 44.71, 128.34, 129.00, 131.94, 132.06, 132.60, 132.89, 140.15, 140.21, 142.50, 142.71.

**Compound 16** was prepared from **4** and 4-bromobutyronitrile as described above for **Compound 2** in 94% yield.

<sup>1</sup>H NMR(CDCl<sub>3</sub>): δ 0.97 (t, J = 7.12Hz, 3H, CH<sub>3</sub>), 1.40 (m, 4H, 2CH<sub>2</sub>), 1.85 (Pent., m, 2H, CH<sub>2</sub>), 2.27 (t, J = 7.17Hz, 2H CH<sub>2</sub>CN), 2.30 (s, 6H, 2CH<sub>3</sub>), 2.57 (s, 6H, 2CH<sub>3</sub>), 2.58 (s, 6H, 2CH<sub>3</sub>), 3.13 (m, 6H, NCH<sub>2</sub>), 3.28 (t, J = 7.11Hz, 2H, NCH<sub>2</sub>), 6.94 (s, 2H, Ph), 6.96 (s, 2H, Ph); <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ 12.55, 14.54, 20.84, 22.64, 22.73, 23.65, 24.43, 24.57, 39.88, 44.31, 44.54, 45.58, 118.69, 131.84, 132.05, 132.73, 133.36, 139.94, 142.20, 142.71.

**Compound 17** was prepared from **16** as described above for **Compound 3** in 93% yield.

<sup>1</sup>H NMR(CDCl<sub>3</sub>): δ 1.00 (t, J = 6.92Hz, 3H, CH<sub>3</sub>), 1.40 (m, 10H, 4CH<sub>2</sub>, NH<sub>2</sub>), 2.29 (s, 6H, 2CH<sub>3</sub>), 2.57 (b, 14H, 4CH<sub>3</sub>, CH<sub>2</sub>N), 3.13 (m, 8H, 4CH<sub>2</sub>N), 6.93 (s, 4H, 2 Ph); <sup>13</sup>C NMR (CDCl<sub>3</sub>): 12.72, 20.90, 22.72, 22.78, 24.67, 24.80, 30.80, 40.02, 41.61, 44.56, 45.10, 45.38, 131.87, 140.04, 142.21, 142.28; MS-FAB(M/Z) 552.3(M<sup>+</sup>, 100%), 368.2, 299.1, 183.0, 154.0.

**Compound 18** was prepared from **17** as described above for **Compound 4**.

<sup>1</sup>H NMR(CDCl<sub>3</sub>): δ 0.96 (t, J = 7.13Hz, 3H, CH<sub>3</sub>), 1.38 (m, 8H, 4CH<sub>2</sub>), 2.29 (s, 9H, 3CH<sub>3</sub>), 2.55 (s, 6H, 2CH<sub>3</sub>), 2.56 (s, 6H, 2CH<sub>3</sub>); 2.59 (s, 6H, 2CH<sub>3</sub>), 2.80 (m, 2H, CH<sub>2</sub>N), 3.10 (m, 8H, NCH<sub>2</sub>), 4.67(t, J = 6.6Hz, 1H, NH), 6.93 (s, 6H, 3 Ph); <sup>13</sup>C NMR(CDCl<sub>3</sub>): δ 12.56, 20.87, 22.70, 22.74, 22.84, 24.40, 26.45, 24.67, 26.62, 39.87, 41.88, 44.45, 45.02,

45.09, 131.86, 131.90, 131.92, 133.12, 133.32, 133.68, 138.91, 139.97, 142.02, 142.21, 142.38; MS-FAB(M/Z): 756.9(M + 23(Na), 100%) 572.8, 390.7, 333.6, 305.6

**Compound 19** was prepared from **4** and 1,4-dichloro-2-butene as described above for **15** in 99% yield. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.01 (t, J=7.11 Hz, 3H, CH<sub>3</sub>), 1.38 (m, 4H, CH<sub>2</sub>), 2.29 (s, 3H), 2.30 (s, 3H), 2.57 (s, 6H), 2.61 (s, 6H), 3.11 (m, 4H, NCH<sub>2</sub>), 3.16 (q, J=7.15 Hz, 2H, NCH<sub>2</sub>), 3.81 (d, J=7.17 Hz, 2H, NCH<sub>2</sub>), 3.98 (d, J=8.05 Hz, 2H, CH<sub>2</sub>Cl), 5.51 (m, 1H, CH=CH), 5.77 (m, 1H, CH-CH), 6.93 (s, 2H, Ph), 6.95 (s, 2H, Ph); <sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 12.76, 20.91, 22.71, 22.76, 24.74, 38.12, 40.08, 41.85, 44.59, 45.54, 129.14, 129.25, 131.88, 132.02, 140.09, 140.19, 142.21, 142.63. MS-FAB (m/z) 569.3 (M<sup>+</sup>, 20%), 385.2, 240.1, 203.3, 183.0, 119 (100%).

**Compound 20** was prepared from **18** and **15** following the procedure described above for **15**. It was purified by column chromatography using hexanes - ethyl acetate (7:3) as eluant (78% yield). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.97 (t, J=7.10 Hz, 3H, CH<sub>3</sub>), 0.99 (t, J=7.0 Hz, 3H, CH<sub>3</sub>), 1.29 (m, 8H, CH<sub>2</sub>), 2.29 (s, 15H, CH<sub>3</sub>), 2.54, 2.55, 2.59 (s, 30H, CH<sub>3</sub>), 3.06 (m, 12H, NCH<sub>2</sub>), 3.65 (m, 8H, NCH<sub>2</sub>), 5.48 (m, 4H, CH=CH), 6.92 (s, 10H, Ph); <sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 12.70, 12.83, 20.88, 20.91, 22.65, 22.68, 22.72, 22.74, 24.48, 24.72, 40.04, 40.47, 41.53, 42.07, 42.22, 42.34, 44.54, 44.96, 127.94, 128.27, 128.57, 129.20, 131.92, 132.05, 139.96, 140.00, 140.12, 140.16, 140.27, 142.19, 142.25, 142.47, 142.58, 142.87. MS-FAB (m/z) 1263.81 (M<sup>+</sup>, 100%), 1080.01, 898.11, 714.81, 563.

**Compound 21:** Pentamide **20** (0.93 g, 0.735 mmol) was dissolved in 20 ml anhydrous CH<sub>2</sub>Cl<sub>2</sub>, phenol (3.46 g, 36.77 mmol) was added, followed by HBr in acetic acid (30%, 17.62 ml) and the mixture was stirred over night at 25°C. Water (10 ml) was added to the flask, the aqueous layer was separated, the organic layer was extracted with 5 ml H<sub>2</sub>O, and the combined aqueous layers were washed with CH<sub>2</sub>Cl<sub>2</sub> (6 x 15 ml). Water was evaporated under vacuum to afford a solid which was dissolved in 1 ml 1N NaOH followed by 1 ml of 50% KOH. This solution was extracted with CHCl<sub>3</sub> (10 x 5 ml). The combined organic layers were dried (MgSO<sub>4</sub>), CHCl<sub>3</sub> was evaporated, and the residue dissolved in anhydrous diethyl ether. Anhydrous HCl gas was passed into the solution while cooling at 0°C. A white solid precipitated which was filtered and washed with ether. It was **21** (84%). <sup>1</sup>H-NMR (D<sub>2</sub>O): δ 1.29 (t, J=7.32 Hz, 3H, CH<sub>3</sub>), 1.31 (t, J=7.24 Hz, 3H, CH<sub>3</sub>), 1.79 (m, 8H, CH<sub>2</sub>), 3.12 (m, 12H, NCH<sub>2</sub>), 3.87 (m, 8H, NCH<sub>2</sub>), 5.98 (m, 4H, CH=CH); <sup>13</sup>C-NMR (D<sub>2</sub>O):



$\delta$  13.36, 13.46, 25.66, 25.77, 45.44, 45.74, 46.24, 46.41, 46.84, 49.09, 49.41, 49.70, 129.02, 129.16, 129.47, 129.66. MS-MALDI (m/z) 354.36 ( $MH^+$ , 100%).

**Compound 22** was prepared in 51% yield from **18** and **14** as described above for compound **15**.  $^1H$ -NMR ( $CDCl_3$ ):  $\delta$  0.97 (t,  $J=6.59$  Hz, 3H,  $CH_3$ ), 0.99 (t,  $J=7.02$  Hz, 3H,  $CH_3$ ), 1.29 (m, 12H,  $CH_2$ ), 2.29 (s, 15H,  $CH_3$ ), 2.55 (s), 2.56 (s), 2.57 (s), 3.10 (m, 16H,  $NCH_2$ ), 3.70 (m, 4H,  $NCH_2$ ), 5.47 (m, 2H,  $CH=CH$ ), 6.93 (s, 10H, Ph);  $^{13}C$ -NMR ( $CDCl_3$ ):  $\delta$  12.69, 12.83, 20.91, 22.69, 22.71, 22.76, 24.43, 24.70, 40.48, 41.11, 41.48, 44.50, 44.91, 128.13, 128.90, 131.88, 131.94, 132.01, 133.29, 139.95, 140.00, 140.15, 142.22, 142.29, 142.60. MS-FAB (m/z) 1265.91 ( $M^+$ , 100%), 1082.01, 900.11, 716.91, 563.81.

**Compound 23** was prepared from **22** in 79% yield as described above for **21**.  $^1H$ -NMR ( $D_2O$ ):  $\delta$  1.29 (t,  $J=7.29$  Hz, 3H,  $CH_3$ ), 1.30 (t,  $J=7.30$  Hz, 3H,  $CH_3$ ), 1.78 (m, 12H,  $CH_2$ ), 3.12 (m, 16H,  $NCH_2$ ), 3.83 (m, 4H,  $NCH_2$ ), 5.96 (m, 2H,  $CH=CH$ );  $^{13}C$ -NMR ( $D_2O$ ):  $\delta$  13.31, 13.42, 25.62, 25.75, 45.38, 45.71, 46.18, 46.76, 49.07, 49.32, 49.69, 129.11, 129.39. MS-MALDI (m/z) 356.38 ( $MH^+$ , 100%).

**Compound 24** was prepared from **18** (52% yield) as described.  $^1H$ -NMR ( $CDCl_3$ ):  $\delta$  0.95 (m, 6H,  $2CH_3$ ), 1.32 (m, 12H,  $CH_2$ ), 2.29 (s, 15H,  $CH_3$ ), 2.55 (s, 30H,  $CH_3$ ), 3.06 (m, 16H,  $NCH_2$ ), 3.70 (m, 4H,  $NCH_2$ ), 5.47 (m, 2H,  $CH=CH$ ), 6.92 (s, 10H, Ph);  $^{13}C$ -NMR ( $CDCl_3$ ):  $\delta$  12.67, 20.90, 22.71, 22.76, 24.43, 24.68, 39.97, 42.08, 44.48, 44.90, 45.61, 128.28, 128.45, 131.87, 131.93, 132.01, 139.96, 140.00, 140.12, 142.21, 142.28, 142.58. MS-FAB (m/z) 1265.91 ( $M^+$ , 100%), 1082.01, 900.11.

**Compound 25** was prepared from **24** in 96% yield as described above for **21**.  $^1H$ -NMR ( $D_2O$ ):  $\delta$  1.28 (t,  $J=7.29$  Hz, 6H,  $2CH_3$ ), 1.78 (m, 12H,  $CH_2$ ), 3.09 (m, 16H,  $NCH_2$ ), 3.84 (m, 4H,  $NCH_2$ ), 5.96 (m, 2H,  $CH=CH$ );  $^{13}C$ -NMR ( $D_2O$ ):  $\delta$  13.31, 25.61, 25.73, 45.70, 46.79, 49.05, 49.36, 49.65, 129.19. MS-MALDI (m/z) 356.4 ( $MH^+$ ).

**Compound 26:** A mixture of KOH (0.25 g),  $K_2CO_3$  (0.25 g) and tetra-*n*-butyl-ammonium hydrogen bromide (0.05 g) were suspended in 15 ml benzene. Mesitylenesulfonylamide (0.199 g, 1 mmol) was added to the suspension and the mixture was heated to 50°C. Iodide **14** (1.98 g, 3 mmol) in 10 ml benzene was added to the flask, the mixture heated under reflux over night, then cooled to room temperature; the inorganic solids were filtered off and washed with benzene (2 x 20 ml). The combined organic layers were washed several times with water until the washings were neutral. The benzene was dried ( $MgSO_4$ ),

evaporated and the residue purified by column chromatography using hexanes and ethyl acetate (7.5:2.5) as eluant; 25% yield (0.948 g).  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  1.00 (t,  $J=7.18$  Hz, 6H,  $\text{CH}_3$ ), 1.28 (m, 8H,  $\text{CH}_2$ ), 2.29 (s, 15H,  $\text{CH}_3$ ), 2.53 (s), 2.55 (s), 2.57 (s), 3.03 (m, 8H,  $\text{NCH}_2$ ), 3.12 (q,  $J=7.13$  Hz, 4H,  $\text{NCH}_2$ ), 3.70 (m, 8H,  $\text{NCH}_2$ ), 5.47 (m, 4H,  $\text{CH}=\text{CH}$ ), 6.93 (s, 10H, Ph);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  12.78, 20.85, 22.63, 22.69, 24.32, 24.58, 40.41, 41.43, 42.00, 44.76, 45.43, 128.08, 128.83, 131.88, 131.95, 132.77, 132.85, 133.23, 139.90, 140.04, 140.08, 142.22, 142.43, 142.53. MS-FAB ( $m/z$ ) 1263.81 ( $\text{M}^+$ , 100%), 1081, 898.11, 815.01, 561.81, 418.81.

**Compound 27** was prepared from **26** in 57% yield as described above for **21**.  $^1\text{H-NMR}$  ( $\text{D}_2\text{O}$ ):  $\delta$  1.31 (t,  $J=7.31$  Hz, 6H,  $\text{CH}_3$ ), 1.78 (m, 8H,  $\text{CH}_2$ ), 3.15 (m, 12H,  $\text{NCH}_2$ ), 3.83 (m, 8H,  $\text{NCH}_2$ ), 5.96 (m, 4H,  $\text{CH}=\text{CH}$ );  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  13.43, 25.64, 25.76, 45.39, 46.19, 46.77, 49.35, 49.72, 129.11, 129.41. MS-MALDI ( $m/z$ ) 354.3 ( $\text{MH}^+$ , 100%).

**Compound 28** was prepared from **15** and mesitylenesulfonylamide in 24% yield as described above for **26**; mp  $57.7^\circ\text{C}$ .  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  0.99 (t,  $J=7.09$  Hz, 6H,  $\text{CH}_3$ ), 2.29 (s, 15H,  $\text{CH}_3$ ), 2.53 (s), 2.55 (s), 3.12 (q,  $J=7.09$  Hz, 4H,  $\text{NCH}_2$ ), 3.63 (m, 16H,  $\text{NCH}_2$ ), 5.49 (m, 8H,  $\text{CH}=\text{CH}$ ), 6.93 (s, 10H, Ph);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  12.85, 20.89, 20.92, 22.66, 40.47, 41.53, 42.19, 128.00, 128.47, 128.58, 129.11, 131.92, 132.05, 140.17, 140.30, 142.46, 142.87. MS-FAB ( $m/z$ ) 1259.81 ( $\text{M}^+$ , 60%), 1075.91, 894.01, 306.51, 153.4 (100%).

**Compound 29** was prepared from **28** in 81% yield as described above for **21**.  $^1\text{H-NMR}$  ( $\text{D}_2\text{O}$ ):  $\delta$  1.31 (t,  $J=7.29$  Hz, 6H,  $\text{CH}_3$ ), 3.15 (q,  $J=7.31$  Hz, 4H,  $\text{NCH}_2$ ), 3.84 (m, 4H,  $\text{NCH}_2$ ), 3.90 (m, 12H,  $\text{NCH}_2$ ), 5.98 (m, 8H,  $\text{CH}=\text{CH}$ );  $^{13}\text{C-NMR}$  ( $\text{D}_2\text{O}$ ):  $\delta$  13.42, 45.41, 46.22, 46.44, 129.07, 129.37, 129.42, 129.58. MS-MALDI ( $m/z$ ) 350.31 ( $\text{MH}^+$ ).

**Compound 30** was prepared from **19** in 25% yield as described above for **26**; mp  $62.3^\circ\text{C}$ .  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  0.95 (s,  $J=7.17$  Hz, 6H,  $\text{CH}_3$ ), 1.33 (m, 8H,  $\text{CH}_2$ ), 2.29 (s, 15H,  $\text{CH}_3$ ), 2.54 (s), 2.55 (s), 3.07 (m, 12H,  $\text{NCH}_2$ ), 3.65 (m, 8H,  $\text{NCH}_2$ ), 5.48 (m, 4H,  $\text{CH}=\text{CH}$ ), 6.92 (s, 10H, Ph);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  12.69, 20.90, 22.69, 22.73, 24.70, 40.03, 42.13, 42.30, 44.53, 45.59, 128.11, 128.79, 131.87, 132.00, 140.02, 140.14, 140.28, 142.17, 142.58, 142.85. MS-FAB ( $m/z$ ) 1263.81 ( $\text{M}^+$ , 100%), 1080.01, 898.11, 714.01, 153.

**Compound 31** was prepared from **30** in 87% yield as described above for **21**.  $^1\text{H-NMR}$  ( $\text{D}_2\text{O}$ ):  $\delta$  1.28 (t,  $J=7.32$  Hz, 6H,  $\text{CH}_3$ ), 1.79 (m, 8H,  $\text{CH}_2$ ), 3.10 (m, 12H,  $\text{NCH}_2$ ), 3.87 (m,

8H, NCH<sub>2</sub>), 5.98 (m, 4H, CH=CH), <sup>13</sup>C-NMR (D<sub>2</sub>O): δ 12.70, 25.00, 25.13, 45.10, 45.81, 46.21, 48.44, 48.78, 128.44, 128.85. MS-MALDI (m/z) 354.3 (MH<sup>+</sup>).

**Compound 32:** Mesitylenesulfonylamide (1.47 g, 7.38 mmol) was dissolved in 50 ml anhydrous DMF, and NaH (85%, 0.3 g) was added to it under a nitrogen atmosphere. The mixture was stirred at room temperature and **19** (1.40 g, 2.46 mmol) in 25 ml DMF were added. Heating at 65°C continued over night. The mixture was cooled to room temperature, and 10 ml of H<sub>2</sub>O were added. The solvents were evaporated and the solid residue was partitioned between 40 ml H<sub>2</sub>O and 40 ml CHCl<sub>3</sub>. The aqueous layer was extracted with CHCl<sub>3</sub> (2 x 30 ml), the combined organic layers were washed with H<sub>2</sub>O (3 x 50 ml), dried (MgSO<sub>4</sub>), and evaporated. The residue was purified by column chromatography using hexanes - ethyl acetate (7.5:2.5). 1.7 g (97%) of **32** as a white solid was obtained. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.94 (t, J=7.10 Hz, 3H, CH<sub>3</sub>), 1.30 (m, 4H, CH<sub>2</sub>), 2.29 (s), 2.30 (s), 2.55 (s, 12H, CH<sub>3</sub>), 2.65 (s, 6H, CH<sub>3</sub>), 3.11 (m, 6H, NCH<sub>2</sub>), 3.52 (m, 1H, NCH), 3.65 (m, 2H, NCH<sub>2</sub>), 3.71 (m, 1H, NCH<sub>2</sub>), 4.82 (br, 1H, NH), 5.47 (m, 2H, CH=CH), 6.93 (s, 4H, Ph), 6.96 (s, 2H, Ph); <sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 12.50, 20.91, 22.71, 22.76, 22.83, 22.91, 24.66, 38.98, 39.85, 42.15, 42.26, 44.50, 128.06, 128.51, 131.86, 131.91, 138.18, 140.00, 140.14, 140.28, 142.17, 142.65.

**Compound 33** was prepared from **32** and **14** in 51% yield as described above for **22**. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.99 (t, J=7.19 Hz, 6H, CH<sub>3</sub>), 1.33 (m, 8H, CH<sub>2</sub>), 2.29 (s, 15H, CH<sub>3</sub>), 2.55 (s), 2.57 (s), 3.10 (m, 12H, NCH<sub>2</sub>), 3.70 (m, 4H, NCH<sub>2</sub>), 3.77 (m, 4H, NCH<sub>2</sub>), 5.42 (m, 4H, CH=CH), 6.93 (s, 10H, Ph); <sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 12.70, 12.71, 20.89, 22.66, 22.72, 22.78, 22.81, 24.60, 26.53, 40.39, 41.37, 41.87, 42.20, 45.47, 128.26, 128.62, 131.78, 131.84, 131.86, 131.92, 132.77, 138.92, 139.96, 140.09, 140.17, 142.57, 142.63.

**Compound 34** was prepared from **33** as described above for **21** in 40% yield.

**Compound 35** was prepared from **15** in 94% yield as described above for **32**.

**Compound 36** was prepared from **35** and **14** in 82% yield as described above for **33**. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.99 (t, J=7.11 Hz, 6H, CH<sub>3</sub>), 1.33 (m, 4H, CH<sub>2</sub>), 2.29 (s, 15H, CH<sub>3</sub>), 2.55 (s), 2.57 (s), 3.07 (m, 8H, NCH<sub>2</sub>), 3.70 (m, 12H, NCH<sub>2</sub>), 5.46 (m, 6H, CH=CH), 6.92 (s, 10H, Ph); <sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 12.69, 12.80, 20.84, 22.62, 22.68, 22.73, 22.77, 24.58, 26.55, 40.44, 41.51, 41.86, 42.04, 42.24, 45.49, 128.10, 128.25, 128.52, 128.62, 128.82,

131.89, 131.95, 132.79, 138.89, 140.07, 140.14, 140.23, 141.94, 142.44, 142.53, 142.82. MS-FAB (m/z) 1262.8 ( $M^+$ , 75%), 1080.01, 896, 119 (100%).

**Compound 37** was prepared from **36** in 65% yield as described above for **21**.  $^1\text{H-NMR}$  ( $\text{D}_2\text{O}$ ):  $\delta$  1.31 (t,  $J=6.97$  Hz, 6H,  $\text{CH}_3$ ), 1.79 (m, 4H,  $\text{CH}_2$ ), 3.12 (m, 8H,  $\text{NCH}_2$ ), 3.83 (m, 12H,  $\text{NCH}_2$ ), 5.96 (m, 6H,  $\text{CH}=\text{CH}$ );  $^{13}\text{C-NMR}$  ( $\text{D}_2\text{O}$ ):  $\delta$  13.48, 25.69, 26.76, 41.67, 45.44, 46.24, 46.45, 46.80, 49.41, 129.00, 129.12, 129.45, 129.71. MS-MALDI (m/z) 352.3 ( $\text{MH}^+$ ).

**Compound 38** was prepared from **35** and **19** in 89% yield as described.  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  0.95 (m, 6H,  $\text{CH}_3$ ), 1.33 (m, 4H,  $\text{CH}_2$ ), 2.29 (s, 15H,  $\text{CH}_3$ ), 2.54 (s), 2.55 (s), 2.57 (s), 3.09 (m, 8H,  $\text{NCH}_2$ ), 3.66 (m, 12H,  $\text{NCH}_2$ ), 5.48 (m, 6H,  $\text{CH}=\text{CH}$ ), 6.93 (s, 10H, Ph);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  12.51, 12.63, 20.84, 20.86, 22.63, 22.65, 22.84, 24.61, 38.92, 40.40, 41.40, 42.11, 42.18, 44.44, 45.48, 127.95, 128.07, 128.49, 128.62, 128.80, 131.76, 131.83, 131.85, 131.88, 132.01, 138.05, 139.01, 140.07, 140.13, 140.24, 142.15, 142.21, 142.87. MS-FAB (m/z) 1263.1 ( $M^+$ , 90%), 1080.1, 896.01, 119 (100%).

**Compound 39** was prepared from **38** in 54% yield as described above for **21**; mp  $270^\circ\text{C}$  (dec.).  $^1\text{H-NMR}$  ( $\text{D}_2\text{O}$ ):  $\delta$  1.31 (m, 6H,  $\text{CH}_3$ ), 1.80 (m, 4H,  $\text{CH}_2$ ), 3.10 (m, 8H,  $\text{NCH}_2$ ), 3.86 (m, 12H,  $\text{NCH}_2$ ), 5.98 (m, 6H,  $\text{CH}=\text{CH}$ );  $^{13}\text{C-NMR}$  ( $\text{D}_2\text{O}$ ):  $\delta$  13.30, 13.42, 25.58, 25.70, 45.69, 46.21, 46.43, 46.81, 49.02, 49.37, 129.00, 129.15, 129.37, 129.59. MS-MALDI (m/z): 352.343 ( $\text{MH}^+$ ).

**Compound 42**: NaH (80%, 132 mg, 4.4 mmol) was added to a solution of diamide **41** (1.98 g, 4.4 mmol) in DMF (10 ml). The mixture was stirred at  $20^\circ\text{C}$  for 30 minutes and a solution of the diester **40** (Reddy et al. (1998) J. Med Chem., 41:4723) (960 mg, 2 mmol) in DMF (10 ml) was added dropwise. The mixture was stirred at  $75^\circ\text{C}$  for 2 h, the solvent was distilled off, the residue was taken in chloroform, washed with a saturated solution of ammonium chloride, dried ( $\text{Na}_2\text{SO}_4$ ) and evaporated to dryness. The crude oil was purified by column chromatography using hexane - ethyl acetate (8:2) as running solvent. 1.4 g (70%) was obtained as a glassy oil.  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  20.58, 22.63, 22.80, 32.42, 33.86, 43.16, 45.42, 46.26, 132.75, 133.21, 139.82, 142.40. MS-FAB 984 ( $M^+$ ),

**Compound 43**: Phenol (1.86 g, 19.7 mmol) and 30% HBr in glacial acetic acid (35 ml) were added in that order to a solution of **42** (600 mg, 0.6 mmol) in  $\text{CH}_2\text{Cl}_2$  (35 ml) at room temperature. The solution was stirred for 24 h, water (30 ml) was added, followed by extraction with methylene chloride (3 x 20 ml). The aqueous layer was evaporated under

reduced pressure and the residue was taken up in 2N NaOH (2 ml) and then 50% KOH (2 ml) followed by extraction with chloroform (6 x 10 ml). After removal of chloroform, the residue was taken up in ethanol (15 ml) and acidified with concentrated hydrochloric acid (0.4 ml). The product **43** (230 mg, 93%) was recrystallized from aqueous ethanol; mp  
5 >270°C (decomp). <sup>1</sup>H-NMR (D<sub>2</sub>O): δ 1.95 (m, 2H), 2.05-2.25 (m, 6H), 2.75 (s, 6H), 2.90 (b, 2H), 3.10-3.35 (m, 12H); <sup>13</sup>C-NMR (D<sub>2</sub>O): δ 25.21, 25.24, 35.60, 35.64, 47.41, 48.58, 50.87. MS-MALDI (m/z) 240 (M<sup>+</sup> + 1).

**Compound 47:** NaH (80%, 132 mg, 4.4 mmol) was added to a solution of diamide **46** (1.98 g, 4.4 mmol) in DMF (10 ml). The mixture was stirred at 20°C for 30 min and a  
10 solution of the diester **8** (900 mg, 2 mmol) in DMF (10 ml) was added dropwise. The mixture was stirred at 75°C for 2 h. The solvent was distilled off, the residue was taken up in chloroform, washed with a saturated solution of ammonium chloride, dried (NaSO<sub>4</sub>) and evaporated to dryness. The oily residue was crystallized from ethyl acetate/hexane 1.2 g (61%); mp 165-166°C. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.08 (t, 3H), 1.75 (m, 4H), 2.28 (s, 12H), 2.55  
15 (bs, 24H), 3.10 (m, 12H), 3.98 (s, 4H), 6.95 (m, 8H); <sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 12.70, 20.86, 22.64, 25.14, 34.85, 40.22, 42.62, 43.37, 78.80, 131.99, 132.26, 133.21, 140.26, 142.28, 142.71. MS-FAB (m/z) 982 (M<sup>+</sup>).

**Compound 48** was obtained as described for **47**. From 1.2 g (1.22 mmol) of tetramide **47**, 420 mg (86%) of the tetrahydrochloride **48** was obtained; mp >270°C (decomp). <sup>1</sup>H-NMR  
20 (D<sub>2</sub>O): δ 1.29 (t, 6H), 2.13 (m, 4H), 3.14 (m, 12H), 4.06 (s, 4H); <sup>13</sup>C-NMR (D<sub>2</sub>O): δ 13.34, 25.52, 39.45, 45.90, 45.64, 46.71, 81.32. MS-MALDI (m/z) 255 (M<sup>+</sup> + 1).

**Compound 44** was obtained as described for **47**. From 450 mg (1 mmol) of diester **8** and 994 mg (2.2 mmol) of diamide **41**, 500 mg (52%) of the tetramide **44** was obtained and crystallized from ethyl acetate - hexane; mp 155-156°C.

25 **Compound 45** was obtained as described for **43**. From 500 mg (0.52 mmol) of tetramide **44**, 160 mg (82%) of the tetrahydrochloride **45** was obtained; mp >270°C (decomp). <sup>1</sup>H-NMR (D<sub>2</sub>O): δ 2.15 (m, 4H), 2.73 (s, 3H), 3.05-3.40 (m, 8H), 4.10 (s, 4H); <sup>13</sup>C-NMR (D<sub>2</sub>O): δ 25.59, 35.66, 45.90, 46.57, 48.61.

**Compound 51** is a mixture of *cis/trans*- isomers. <sup>1</sup>H-NMR (D<sub>2</sub>O): δ 1.15-2.10 (m, 7H),  
30 2.90 (q, 1H), 3.30-3.80 (b, 2H); <sup>13</sup>C-NMR (D<sub>2</sub>O): δ 24.16, 24.97, 28.44, 30.42, 36.58, 37.14, 48.24, 52.27, 55.19, 57.45, 64.55, 67.26.

**Compound 52:** Mesitylenesulfonylchloride (6.5 g, 30 mmol) in dioxane (10 ml) was added dropwise to a stirred and cooled mixture of amine alcohol **51** (1.15 g, 10 mmol), triethylbenzyl ammonium bromide (135 mg, 0.5 mmol), 50% KOH (10 ml) and dioxane (10 ml). The reaction mixture was left over night at 20 °C with magnetic stirring. An excess of water was added, the solution was extracted with chloroform (3 x 30 ml), dried ( $\text{Na}_2\text{SO}_4$ ) and evaporated to dryness. The oily residue was chromatographed on a silica-gel column using hexane:ethyl acetate (8:2) as eluants. Crystallization from ethyl acetate - hexane afforded 1.2 g (25%) of pure **52**; mp 167-168°C.  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  1.35-1.90 (6H), 1.90-2.15 (m, 1H), 2.30, 2.35 (s, 6H), 2.65 (s, 12H), 3.20 (m, 1H), 3.70 (m, 1H), 3.90 (m, 1H), 5.15 (d, 1H), 6.90, 7.00 (s, 4H);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  20.73, 20.85, 22.15, 22.37, 22.70, 26.94, 32.75, 45.34, 56.09, 70.38, 130.22, 131.57, 133.98, 138.68, 139.64, 142.02, 143.10. MS-EI (m/z) 479 ( $\text{M}^+$ ), 280 ( $\text{M}^+ - 199$ ).

**Compound 54:** NaH (105 mg, 3.5 mmol) was added to a solution of compound **52** (1.7 g, 3.5 mmol) in DMF (10 ml). The mixture was stirred at 20°C for 30 min and a solution of compound **53** (1.34 g, 3.85 mmol) in DMF (5 ml) was added in small portions. The mixture was stirred at 75 °C for 2 h. The solvent was distilled off, the residue was taken up in chloroform, washed with a saturated solution of ammonium chloride, dried ( $\text{Na}_2\text{SO}_4$ ) and evaporated. The oily residue was purified by column chromatography (hexane - ethyl acetate 8:2) which gave compound **54** (1.22 g, 47%).  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  1.98 (t, 3H), 1.20-2.05 (9H), 2.20 (s, 6H), 2.55, 2.65 (s, 12H), 2.70-3.55 (9H), 6.85 (s, 4H);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  12.49, 20.80, 21.64, 21.87, 22.88, 28.72, 33.16, 36.13, 39.96, 43.80, 47.95, 57.77, 61.26, 131.83, 132.94, 133.14, 138.82, 139.90, 142.07, 142.63. MS-FAB (m/z) 628 ( $\text{M}^+ + 1$ ), 546 ( $\text{M}^+ - 81$ ).

**Compound 55** was obtained following the procedure described for compound **42**. From 1.22 g (1.6 mmol) of bromoderivative **54** and 820 mg (1.76 mmol) of diamide **46**, 1.26 g (77%) of tetramide **55** was obtained as a glassy oil.  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  0.80 (t, 6H), 1.20-1.75 (6H), 1.90 (m, 1H), 2.15 (s, 12H), 2.35-2.60 (s, 24H), 2.65-3.40 (15H), 6.85 (b, 8H);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  12.38, 20.71, 22.52, 22.66, 24.72, 27.55, 28.04, 39.19, 39.71, 41.02, 42.33, 42.62, 43.37, 48.81, 61.44, 131.76, 131.88, 133.10, 133.89, 138.66, 139.93, 142.17, 142.33, 142.57. MS-FAB (m/z) 1012 ( $\text{M}^+$ ), 828 ( $\text{M}^+ - 184$ ).

**Compound 56** was obtained following the procedure described for compound 43. From 1.26 g (1.24 mmol) of tetramide 55, 300 mg (56%) of the tetrahydrochloride 56 was obtained; mp > 270°C (decomp). <sup>1</sup>H-NMR (D<sub>2</sub>O): δ 1.35 (t, 6H), 1.60 (m, 1H), 1.80 (b, 3H), 2.15 (b, 6H), 2.50 (b, 1H), 3.20 (m, 13H), 3.45 (m, 2H); <sup>13</sup>C-NMR (D<sub>2</sub>O): δ 13.23, 25.48, 25.73, 25.79, 31.69, 31.99, 43.40, 45.91, 46.43, 46.71, 48.07, 53.20, 75.28. MS-MALDI (m/z) 285 (M<sup>+</sup> + 1).

**Compound 57:** NaH (80%, 150 mg, 5 mmol) and NaBr (2.5 g, 25 mmol) were added to a solution of compound 52 (2.35 g, 4.9 mmol) in DMF (15 ml). The mixture was stirred at 20°C for 30 min and a solution of 1-bromoethane (2.2 g, 25 mmol) in DMF (10 ml) was added in small portions. The mixture was stirred at 90°C for 3 h. The solvent was distilled off, the residue taken up in chloroform, washed with a saturated solution of ammonium chloride, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The product was purified by silica gel chromatography (hexane/ethyl acetate 9:1). The oily residue (1.5 g, 79%) crystallized on standing; mp 68-69°C. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.10 (t, 3H), 1.30-2.10 (6H), 2.25 (b, 4H), 2.60 (s, 6H), 3.20 (m, 2H), 3.35 (m, 2H), 3.60 (m, 2H), 6.95 (s, 2H); <sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 16.35, 20.93, 21.79, 22.89, 29.32, 29.37, 36.54, 38.12, 44.13, 61.40, 131.99, 132.80, 140.20, 142.52. MS-FAB 389 (M<sup>+</sup> + 1), 308 (M<sup>+</sup> - 80).

**Compound 59** was obtained following the procedure described for compound 42. From 700 mg (1.80 mmol) of compound 57 and 394 mg (0.9 mmol) of diamide 58, 400 mg (37%) of tetramide 59 were obtained. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.90 (t, 6H), 1.25-1.80 (m, 8H), 1.80-2.10 (m, 8H), 2.15 (s, 12H), 2.40, 2.50 (s, 24H), 2.60-3.35 (m, 6H), 2.85, 2.90 (s, 8H); <sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 16.14, 20.85, 21.95, 21.99, 22.55, 25.49, 28.78, 28.88, 31.49, 37.87, 40.50, 40.83, 43.85, 44.06, 49.30, 61.42, 131.86, 131.96, 133.09, 133.40, 139.93, 139.98, 142.27, 142.40. MS-FAB (m/z) 1052 (M<sup>⊕</sup>), 891 (M<sup>+</sup> - 184).

**Compound 60** was obtained following the procedure described for compound 43. From 400 mg (0.38 mmol) of tetramide 59, 95 mg (53%) of the tetrahydrochloride derivative were obtained; mp > 270°C (decomp.) <sup>1</sup>H-NMR (D<sub>2</sub>O): δ 1.30 (t, 6H), 1.60 (m, 2H), 1.80 (m, 6H), 1.95-2.35 (6H), 2.45 (m, 2H), 3.20 (m, 10H), 3.40 (m, 4H); <sup>13</sup>C-NMR (D<sub>2</sub>O): δ 13.59, 25.34, 25.71, 31.75, 32.00, 43.34, 44.83, 48.02, 53.24, 64.52. MS-MALDI (m/z) 325 (M<sup>+</sup> + 1).

**Compound 62:** Mesitylenesulfonylchloride (3.27 g, 15 mmol) in dioxane (20 ml) was added dropwise to a stirred solution of **61** (1.3 g, 10 mmol) in dioxane (20 ml) and 50% KOH (15 ml) at 0°C. When addition was completed, the mixture was left over night at 20°C. Excess water was added, the solution cooled and the precipitate filtered off.

5 Crystallization from ethylacetate - hexane gave compound **62** (2 g, 80%); mp 115-116°C. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 2.35 (s, 3H), 2.55 (t, 2H), 2.65 (s, 6H), 3.25 (q, 2H), 5.15 (t, 1H), 7.0 (s, 2H); <sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 19.07, 20.82, 22.78, 38.37, 117.56, 132.07, 133.0, 138.99, 142.67. MS-EI (m/z) 252 (M<sup>+</sup>).

**Compound 63:** NaH (80%, 330 mg, 11 mmol) was added to a solution of compound **62** (2.52 g, 10 mmol) in DMF (20 ml) under N<sub>2</sub>. The mixture was stirred for 30 min and a solution of compound **53** (3.82 g, 11 mmol) in DMF (10 ml) was added in small portions. The mixture was stirred at 70°C for 2h. The solvent was distilled off, the residue taken up in chloroform, washed with a saturated solution of ammonium chloride, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated to dryness. The product was purified by silica-gel chromatography (hexane - ethyl acetate 8:2). The oily residue (3.0 g, 57%) crystallized on standing; mp 105-106°C. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.00 (t, 3H), 1.75 (m, 2H), 2.35 (s, 6H), 2.60 (14H), 3.10 (m, 6H), 3.45 (t, 3H), 6.90, 6.95 (s, 4H); <sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 12.63, 16.94, 20.89, 22.67, 25.73, 40.27, 42.19, 42.51, 44.72, 117.36, 131.95, 132.22, 140.06, 140.34, 142.52, 143.33. MS-EI (m/z) 519 (M<sup>+</sup>), 429 (M<sup>+</sup> - HCN).

20 **Compound 65:** The nitrile **63** (3.0 g, 5.7 mmol) was dissolved in a mixture of ethanol (150 ml) and concentrated hydrochloric acid (1.5 ml). PtO<sub>2</sub> was added (300 mg), the mixture was hydrogenated at 50 psi over night, the catalyst was filtered off and the solvent evaporated to afford an oily residue of compound **64**, which was used in the next step without further purification. Free base <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.00 (t, 3H), 1.55 (m, 2H), 1.75 (m, 2H), 2.30 (s, 6H), 2.55 (14 H), 2.90-3.30 (8H), 6.95 (s, 4H); <sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 12.64, 20.87, 22.69, 25.35, 30.93, 39.04, 40.12, 42.65, 43.11, 131.86, 133.10, 140.04, 142.43. MS-FAB (m/z) 524 (M<sup>+</sup> + 1).

30 Mesitylenesulfonylchloride (1.86 g, 8.55 mmol) in dioxane (15 ml) was added dropwise to a stirred mixture of **64** (3.0 g, 5.7 mmol) dissolved in dioxane (30 ml) and 50% KOH (15 ml) at 0°C. The reaction mixture was allowed to reach room temperature and was kept for further 2 h. An excess of water was added and the mixture was extracted with chloroform, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated to dryness. Purification was achieved by silica gel column



chromatography using hexane - ethyl acetate (8:2) as eluant; 2.79 g (69%) of **65** were obtained.  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  0.95 (t, 3H), 1.60 (m, 4H), 2.30 (s, 9H), 2.50 (s, 12H), 2.65 (s, 6H), 2.85 (m, 2H), 3.05 (6H), 3.20 (t, 2H), 5.00 (t, 1H), 6.95 (6H);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  12.45, 20.81, 22.73, 25.23, 27.46, 39.19, 33.99, 42.49, 42.92, 43.17, 131.84, 133.05, 133.82, 138.80, 139.90, 141.92, 142.36, 142.64. MS-FAB ( $m/z$ ) 705 ( $\text{M}^\oplus$ ).

**Compound 66** was obtained following the procedure described for compound **42**. From 705 mg (1 mmol) of **65** and 426 mg (1.1 mmol) of **57**, 470 mg (46%) of tetramide **66** was obtained as a glassy product.  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  0.85-1.10 (t, 6H), 1.35-2.10 (m, 11H), 2.30 (s, 12H), 2.40-2.65 (m, 24H), 2.75-3.55 (m, 13H), 6.95 (m, 8H);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  12.64, 16.11, 20.91, 22.08, 22.75, 24.81, 25.09, 28.83, 29.07, 37.93, 40.08, 40.84, 42.50, 42.81, 43.11, 43.42, 49.11, 61.43. MS-FAB ( $m/z$ ) 1013 ( $\text{M}^+ + 1$ ).

**Compound 67** was obtained following the procedure described for compound **43**. From 470 mg (0.46 mmol) of tetramide **66**, 142 mg (71%) of the tetrahydrochloride derivative was obtained; mp  $>250^\circ\text{C}$  (decomp).  $^1\text{H-NMR}$  ( $\text{D}_2\text{O}$ ):  $\delta$  1.30 (t, 6H), 1.60 (m, 1H), 1.85 (b,s, 3H), 2.15 (m, 6H), 2.45 (m, 1H), 3.15 (m, 13H), 3.45 (m, 2H);  $^{13}\text{C-NMR}$  ( $\text{D}_2\text{O}$ ):  $\delta$  13.29, 13.57, 25.34, 25.44, 25.64, 31.68, 31.94, 43.27, 44.80, 45.86, 46.62, 47.42, 47.97, 53.19, 64.50. MS-MALDI 285 ( $\text{M}^+ + 1$ ), 286 ( $\text{M}^+ + 2$ ).

**Compound 68a**: 4-Cyanobenzaldehyde (Aldrich, 1.31 g, 10 mmol) was dissolved in 30 ml anhydrous MeOH followed by the addition of  $\text{MgSO}_4$  (anhydrous, 1.5 g) and 1,4-diaminobutane (Aldrich, 0.44 g, 5 mmol) and the mixture was stirred under argon over night. The suspension was cooled in an ice bath and  $\text{NaBH}_4$  (2.0 g) was added in portions and stirring continued for 2 h at  $0^\circ\text{C}$ . The methanol was evaporated under vacuum and the resulting solid was partitioned between 35 ml  $\text{H}_2\text{O}$  and 50 ml  $\text{CHCl}_3$ . Some of the solid was not soluble in either the  $\text{H}_2\text{O}$  or the  $\text{CHCl}_3$  and was filtered off and the aqueous layer was extracted with  $\text{CHCl}_3$  (2 x 25 ml). The pooled organic layers were dried ( $\text{MgSO}_4$ ), evaporated and the solid was recrystallized from ethyl acetate - hexane, yield 1.1 g (35%); mp  $90.6-90.8^\circ\text{C}$ .  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  1.43 (broad, 2H, NH), 1.55 (m, 4H,  $\text{CH}_2$ ), 2.63 (m, 4H,  $\text{NCH}_2$ ), 3.85 (s, 4H, benzylic  $\text{CH}_2$ ), 7.44 (m, 4H, Ph), 7.60 (m, 4H, Ph);  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  27.78, 49.28, 53.44, 110.65, 118.88, 128.52, 132.12, 146.21. MS ( $m/z$ ) 318 ( $\text{M}^+$ ), 185, 145, 131, 116 (100%), 70.

**Compound 68b** was prepared from 4-cyano-benzaldehyde and 1,5-diaminopentane as described above for **68a**; 42% yield; mp 92.9-93.0°C. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.40 (m, 4H, NH, CH<sub>2</sub>), 1.50 (m, 4H, CH<sub>2</sub>), 2.59 (m, 4H, NCH<sub>2</sub>), 3.83 (s, 4H, benzylic CH<sub>2</sub>), 7.45 (m, 4H, Ph), 7.59 (m, 4H, Ph); <sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 24.86, 29.87, 49.29, 53.40, 110.50, 118.85, 128.48, 132.04, 146.19. MS (m/z) 332 (M<sup>+</sup>), 216, 199, 145, 116 (100%), 84.

**Compound 68c** was prepared from 4-cyanobenzyldehyde and 1,6-diaminohexane as described above for **68a**; 45% yield; mp 95.6-95.8°C. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.35 (m, 4H, CH<sub>2</sub>), 1.50 (m, 6H, NH, CH<sub>2</sub>), 2.60 (t, J=6.92 Hz, 4H, NCH<sub>2</sub>), 3.84 (s, 4H, benzylic CH<sub>2</sub>), 7.44 (m, 4H, Ph), 7.60 (m, 4H, Ph); <sup>13</sup>C-NMR (CDCl<sub>3</sub>): δ 27.17, 30.02, 49.42, 53.50, 110.65, 118.92, 128.55, 132.14, 146.27. MS (m/z) 346 (M<sup>+</sup>), 230, 213, 145, 116 (100%) 98.

**Compound 69a:** Dinitrile **68a** (0.75 g, 2.36 mmol) was dissolved in anhydrous THF, lithium bis(trimethylsilyl)amide (9.43 ml of a 1 M solution in THF) was added slowly under argon atmosphere. The mixture was stirred at room temperature for 2 h; then cooled in an ice bath, followed by the addition of 4 equivalents of 6N HCl in ether. A white solid precipitated immediately and was filtered after 12 h. The solid was recrystallized from ethanol-ether to afford 1.19 g of compound **69a** (93%). <sup>1</sup>H-NMR (D<sub>2</sub>O): δ 1.87 (m, 4H, CH<sub>2</sub>), 3.22 (m, 4H, CH<sub>2</sub>N), 4.40 (s, 4H, benzylic CH<sub>2</sub>), 7.74 (m, 4H, Ph), 7.91 (m, 4H, Ph); <sup>13</sup>C-NMR (DMSO-d<sub>6</sub>): δ 22.68, 46.09, 49.28, 128.10, 128.47, 130.69, 138.15, 165.44. MS-ESI (m/z) 353.2 (M<sup>+</sup>), 177.2 (100%).

**Compound 69b** was prepared from **68b** in 92% yield as described above for **69a**. <sup>1</sup>H-NMR (D<sub>2</sub>O): δ 1.52 (m, 2H, CH<sub>2</sub>), 1.80 (m, 4H, CH<sub>2</sub>), 3.19 (m, 4H, NCH<sub>2</sub>), 4.40 (s, 4H, benzylic CH<sub>2</sub>), 7.75 (m, 4H, Ph), 7.91 (m, 4H, Ph); <sup>13</sup>C-NMR (DMSO-d<sub>6</sub>): δ 24.90, 26.91, 48.96, 51.88, 130.29, 130.46, 132.43, 139.51, 167.52. MS-ESI (m/z) 367.2 (M<sup>+</sup>), 350.2 (100%), 301.2.

**Compound 69c** was prepared from **68c** as described above for **69a** in 96% yield. <sup>1</sup>H-NMR (D<sub>2</sub>O): δ 1.46 (m, 4H, CH<sub>2</sub>), 1.78 (m, 4H, CH<sub>2</sub>), 3.16 (m, 4H, NCH<sub>2</sub>), 4.39 (s, 4H, benzylic CH<sub>2</sub>), 7.74 (m, 4H, Ph), 7.91 (m, 4H, Ph); <sup>13</sup>C-NMR (DMSO-d<sub>6</sub>): δ 25.24, 25.82, 46.73, 49.44, 128.35, 128.56, 130.81, 138.38, 165.58. MS-ESI (m/z) 381.2 (M<sup>+</sup>), 191.2 (100%), 150, 116.

**Compound 70:** Triamide **18** (4.3 g, 5.8 mmol) was dissolved in 30 ml of DMF and 80% NaH (208 mg, 6.9 mmol) was added. The mixture was stirred under a N<sub>2</sub> atmosphere for 1 h and 1.12 g (7.5 mmol) of bromobutyronitrile dissolved in 3 ml of DMF were added all at once. The reaction mixture was heated for 3 h at 90°C. The solvent was distilled-off and the residue was dissolved in chloroform and washed twice with a saturated solution of ammonium chloride; dried (NaSO<sub>4</sub>) and evaporated to dryness. Flash chromatography of the residue using hexane – ethyl acetate (6:4) as eluant gave the yellow oil **70** (3.7 g, 77%).  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.95 (t, 3H), 1.35 (m, 8H), 1.85 (m, 2H), 2.20 (t, 2H), 2.30 (s, 9H), 2.55 (s, 18H), 3.10 (m, 10H), 3.25 (t, 2H), 6.95 (s, 6H). MS-FAB (m/z) 823 (M<sup>+</sup> + Na), 639, 457.

**Compound 71:** Nitrile **70** (3.7 g, 4.6 mmol) was dissolved in 20 ml of chloroform and 150 ml of ethanol were added. The mixture was reduced over 0.35 g of PtO<sub>2</sub> at 50 psi over night. The catalyst was filtered-off and the solvent evaporated to dryness. The oily residue was dried in vacuo for 2 h and dissolved in 50 ml of Cl<sub>3</sub>CH and 12 ml 2N NaOH. The mixture was cooled in an icewater bath with efficient magnetic stirring and 1.50 g (6.9 mmol) of mesitylene chloride dissolved in 10 ml of chloroform were added all at once. After 2 h the organic layer was separated, washed twice with a saturated solution of ammonium chloride, dried (NaSO<sub>4</sub>) and evaporated to dryness. Flash chromatography of the residue using hexane – ethyl acetate (7:3) as eluant provided the tetramide **71** as a colorless oil (3.3 g, 73% over two steps).  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.95 (t, 3H), 1.40 (m, 12H), 2.30 (s, 12H), 2.60 (s, 24H), 2.80 (b, 2H), 3.10 (m, 12H), 4.70 (b, 1H), 6.90 (s, 8H). MS-FAB (m/z) 1010 (M<sup>+</sup> + 1 + Na), 826, 643.

**Compound 72:** The tetramide **71** (6.28 g, 6.3 mmol) was dissolved in 40 ml of DMF and 80% NaH (230 mg, 7.6 mmol) was added. The mixture was stirred under a N<sub>2</sub> atmosphere for 1 h and 1.30 g (8.8 mmol) of bromobutyronitrile dissolved in 3 ml of DMF were added all at once. The reaction mixture was heated for 3h at 90°C, the solvent was distilled-off and the residue was extracted into chloroform and washed twice with a saturated solution of ammonium chloride; dried (NaSO<sub>4</sub>) and evaporated to dryness. Flash chromatography of the residue with hexane – ethyl acetate (7:3) as eluant provided the nitrile **72** (5.0 g, 74%).  
<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.95 (t, 3H), 1.35 (m, 12H), 1.80 (m, 2H), 2.25 (t, 2H), 2.35 (s, 12H), 2.70 (s, 24H), 3.10 (m, 14H), 3.25 (t, 2H), 7.0 (s, 8H). MS-FAB (m/z) 1077 (M<sup>+</sup> + 1 + Na), 893, 711, 586.

**Compound 73:** Nitrile **72** (6.0 g, 5.6 mmol) was dissolved in 20 ml of chloroform and 150 ml of ethanol were added. The mixture was reduced over 600 mg of PtO<sub>2</sub> at 50 psi overnight. The catalyst was filtered-off and the solvent evaporated to dryness. The oily residue was dried in vacuo for 2 h and dissolved in 100 ml of chloroform and 15 ml 2N NaOH. The mixture was cooled in an icewater bath with efficient magnetic stirring, and 1.80 g (8.4 mmol) of mesitylene chloride dissolved in 10 ml of Cl<sub>3</sub>CH was added all at once. After 2h the organic layer was separated, washed twice with a saturated solution of ammonium chloride, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated to dryness. Flash chromatography of the residue using hexane – ethyl acetate (7:3) as eluant gave the pentamide **73** as a colorless oil (5.0 g, 71% over two steps). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.95 (t, 3H), 1.35 (m, 16H), 2.30 (s, 15H), 2.55 (s, 30H), 2.75 (bs, 2H), 3.05 (m, 16H), 4.70 (b, 1H), 6.90 (s, 10H). MS-FAB (m/z) 1261 (M<sup>+</sup> - 1 + Na), 1077, 895.

**Compound 74:** Pentamide **73** (3.4 g, 2.7 mmol) was dissolved in 30 ml of DMF and 60% NaH (162 mg, 4.05 mmol) was added. The mixture was stirred under a N<sub>2</sub> atmosphere for 0.5 h and 2.3 g (10.8 mmol) of 2-bromoethanol benzylether dissolved in 3 ml of DMF were added all at once. The reaction mixture was heated for 2 h at 80°C, the solvent was distilled-off and the residue was dissolved in chloroform and washed twice with a saturated solution of ammonium chloride, dried (NaSO<sub>4</sub>) and evaporated to dryness. Flash chromatography of the residue using hexane – ethyl acetate (7:3) as eluant provided the product **74** (2.6 g, 70%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.95 (t, 3H), 1.30 (m, 16H), 2.30 (s, 15H), 2.50 (s, 30H), 2.90-3.20 (m, 18H), 3.25 (t, 2H), 2.35 (t, 2H), 4.35 (s, 2H), 6.95 (s, 10H), 7.20-7.35 (m, 5H). <sup>13</sup>C NMR (CDCl<sub>3</sub>): δ 12.65, 20.84, 22.67, 22.71, 24.41, 24.66, 39.97, 44.48, 44.88, 46.59, 68.01, 72.95, 127.46, 127.57, 128.25, 131.83, 131.89, 133.28, 139.88, 139.95, 140.04, 142.16, 142.23. MS-FAB (m/z) 1394 (M<sup>+</sup> - 2 + Na) 1030.

**Compound 75:** Pentamide **74** (1.2 g, 0.87 mmol) was dissolved in 12 ml of methylene chloride followed by the addition of 30% HBr/acetic acid (16 ml) and phenol (3.0 g, 32 mmol). The mixture was stirred at room temperature overnight, water (16 ml) was added, followed by extraction with methylene chloride (3 × 10 ml). The aqueous layer was evaporated in vacuo. The residue was dissolved in 2N NaOH (4 ml) and 50% KOH (4 ml) followed by extraction with chloroform (4 × 10 ml). After removal of the solvent the residue was dissolved in ethanol (20 ml) and acidified with concentrated hydrochloric acid (0.5 ml). The white precipitate (**75**) was recrystallized from aqueous ethanol (440 mg,

90%); mp above 270°C (decomp). <sup>1</sup>H-NMR (D<sub>2</sub>O): δ 1.30 (t, 3H), 1.75 (b, 16H), 2.90-3.30 (m, 20H), 2.85 (t, 2H). <sup>13</sup>C NMR (D<sub>2</sub>O): δ 13.29, 25.48, 25.59, 45.70, 49.04, 49.49, 49.67, 51.88, 59.39. MS-MALDI (m/z) 374 (M<sup>+</sup> + 1).

**Compound 76:** Pentamide **73** (850 mg, 0.68 mmol) was dissolved in DMF (15 ml) and 80% NaH (30 mg, 1 mmol) was added. The mixture was stirred under a N<sub>2</sub> atmosphere at room temperature for 0.5 h and 137 mg (0.30 mmol) of **73** dissolved in 5 ml of DMF were slowly added. The reaction mixture was heated for 2 h at 80°C, the solvent was distilled-off and the residue was dissolved in chloroform and washed twice with a saturated solution of ammonium chloride, dried (NaSO<sub>4</sub>) and evaporated to dryness. Flash chromatography of the residue using hexane – ethyl acetate – methanol (6:4:0.1) as eluant afforded the product **76** (590 mg, 77%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.95 (t, 6H), 1.15-1.40 (m, 32H), 2.30 (s, 30H), 2.55 (s, 60H), 2.90-3.25 (m, 36H), 3.60 (d, 4H), 5.40 (t, 2H), 6.95 (s, 20H). MS-FAB 2553 (M<sup>+</sup> + Na).

**Compound 77** was obtained following the procedure described for compound **75**. From 650 mg (0.25 mmol) of decamide **76**, 225 mg (81%) of decahydrochloride **77** was obtained; mp > 270°C (decomp). <sup>1</sup>H-NMR (D<sub>2</sub>O): δ 1.30 (t, 6H), 1.75 (b, 32H), 3.10 (b, 36H), 3.75 (b, 4H), 6.05 (b, 2H); <sup>13</sup>C NMR (D<sub>2</sub>O): δ 13.28, 25.57, 45.66, 49.00, 49.13, 49.64, 50.86, 131.15. MS-ESI 711 (M<sup>+</sup> + 1).

**Compound 78** was obtained following the procedure described for compound **76**. From 850 mg of **73**, 360 mg (47%) of decamide **78** were obtained. <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.95 (t, 6H), 1.15-1.45 (m, 32H), 2.30 (s, 30H), 2.55 (s, 60H), 2.90-3.20 (b, 36H), 3.65 (d, 4H), 5.40 (t, 2H), 6.90 (s, 20H). MS-FAB (m/z) 2553 (M<sup>+</sup> + Na).

**Compound 79** was obtained following the procedure described for compound **75**. From 330 mg (0.13 mmol) of decamide **78**, 127 mg (90%) of decahydrochloride **79** was obtained; mp > 270°C (decomp). <sup>1</sup>H-NMR (D<sub>2</sub>O): δ 1.30 (t, 6H), 1.80 (b, s, 32H), 3.10 (b, 36H), 3.85 (d, 4H), 6.0 (t, 2H). <sup>13</sup>C NMR (D<sub>2</sub>O): δ 13.31, 25.59, 45.71, 46.83, 49.05, 49.39, 49.69, 129.21. MS-ESI (m/z) 512 (M<sup>+</sup> + 2).

**Compound 96:** A suspension of NaH (60% in mineral oil, 336 mg, 14 mmol) in DMF (50 ml) was slowly added to a stirred solution of benzyl-4-bromobuty ether (3.645 g, 15 mmol) and mesitylenesulfonamide **18** (7.33g, 10 mmol) in DMF (180 ml) at 0°C. The reaction mixture was stirred for 10 h at 50°C, quenched with 15 ml of H<sub>2</sub>O at 0°C, acidified to

pH=7 with 5% HCl, extracted with Et<sub>2</sub>O, washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and purified on a column (SiO<sub>2</sub>, EtOAc/Hexane = 3:7); yield 6.9 g, (77%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 0.98 (t, J=7.1 Hz, 3H, CH<sub>3</sub>), 1.25-1.30 (m, 12H, CH<sub>2</sub>), 2.27 (s, 3H, CH<sub>3</sub>), 2.29 (s, 6H, CH<sub>3</sub>), 2.55 (s, 18H, CH<sub>3</sub>), 3.0-3.2 (m, 12H, CH<sub>2</sub>), 3.31 (t, J=6.0 Hz, CH<sub>2</sub>), 4.41 (s, 2H, CH<sub>2</sub>), 6.91 (s, 2H, Ph), 6.92 (s, 4H, Ph), 7.2-7.4 (m, 5H, Ph).

**Compound 97:** A solution of HBr 30% in AcOH (45 ml) was added to a stirred solution of **96** (2 g, 2.24 mmol) and phenole (6.324 g, 67.2 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (23 ml) at 0°C. The cooling bath was removed and the reaction mixture was stirred for 24 h at 20°C. The reaction mixture was quenched with H<sub>2</sub>O (45 ml), washed with CH<sub>2</sub>Cl<sub>2</sub>, and concentrated to dryness *in vacuo*. The residue was cooled to 0°C, basified with 2N NaOH (5 ml), followed by 50% KOH (5 ml). The product was extracted with CHCl<sub>3</sub> (7 x 10 ml); yield 475 mg (81%). <sup>1</sup>H-NMR (D<sub>2</sub>O): δ 1.10 (t, J=7 Hz, 3H, CH<sub>3</sub>), 1.45-1.70 (m, 12H, CH<sub>2</sub>), 2.55-2.70 (m, 12H, CH<sub>2</sub>), 3.57 (t, J=5.0 Hz, CH<sub>2</sub>).

**Compound 98:** A solution of Na<sub>2</sub>CO<sub>3</sub> (10%, 26 ml) was added to triamine **97** (830 mg, 3.20 mmol) in dioxane (21 ml). Di-*tert*-butyl dicarbonate (3.25 g, 24 mmol) solution in dioxane (21 ml) was added into the reaction mixture at 0°C and stirred for 10 h at 20°C. The reaction mixture was diluted with CHCl<sub>3</sub> (200 ml), washed with H<sub>2</sub>O, brine, dried (Na<sub>2</sub>SO<sub>4</sub>), concentrated and purified on a column (SiO<sub>2</sub>, EtOAc/hexane=4:6); yield 1.7 g, (96%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.09 (t, J=7.1 Hz, 3H, CH<sub>3</sub>), 1.45-1.65 (m, 39H, CH<sub>2</sub>, CH<sub>3</sub>), 3.1-3.3 (m, 12H, CH<sub>2</sub>), 3.67 (t, J=6 Hz, CH<sub>2</sub>).

**Compound 99:** Diethyl azodicarboxylate (196 mg, 1.13 mmol) in THF (0.6 ml) was added to a cold mixture of **98** (630 mg, 1.127 mmol), triphenylphosphine (296 mg, 1.13 mmol) and phthalimide (166 mg, 1.13 mmol) in THF (1.2 ml), stirred for 10 h, concentrated *in vacuo*, and purified on a column (SiO<sub>2</sub>, EtOAc/hexane=3:7); yield 835 mg (97%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.09 (t, J=7, 3H, CH<sub>3</sub>), 1.35-1.80 (m, 39H, CH<sub>2</sub>, CH<sub>3</sub>), 3.1-3.35 (m, 12H, CH<sub>2</sub>), 3.70 (t, J=6.7 Hz, 2H, CH<sub>2</sub>), 7.68-7.80 (m, 2H, Phth), 7.80-7.87 (m, 2H, Phth).

**Compound 100:** A mixture of **99** (275 mg, 0.4 mmol), hydrazine monohydrate (98%, 43.5 µl, 0.85 mmol) in EtOH were heated at 80°C for 45 min. The precipitate was filtered and washed with cold EtOH, filtrate was combined with washings, concentrated and dried *in vacuo*; yield 220 mg (99%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ 1.09 (t, J=7.0 Hz, 3H, CH<sub>3</sub>), 1.25-1.57 (m, 39H, CH<sub>2</sub>, CH<sub>3</sub>), 2.71 (t, J=6.7 Hz, 2H, CH<sub>2</sub>), 3.1-3.3 (m, 12H, CH<sub>2</sub>).

**Compound 101** Mesoporphyrin IX dihydrochloride (70 mg, 0.11 mmol) was dissolved in DMF (1 ml), combined with HBTU (83 mg, 0.22 mmol) and diisopropylethyl amine (174  $\mu$ l, 1 mmol). The mixture was stirred for 5 min, combined with **100** (167 mg, 0.3 mmol), stirred for 10 h, quenched with H<sub>2</sub>O, concentrated *in vacuo*, dissolved in CHCl<sub>3</sub>, washed with H<sub>2</sub>O (5 times), 2% KHSO<sub>4</sub>, NaHCO<sub>3</sub> solution, dried (Na<sub>2</sub>SO<sub>4</sub>), and purified by a column chromatography (SiO<sub>2</sub>, MeOH/CHCl<sub>3</sub> = 1:20); yield 177 mg (98%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>):  $\delta$  1.06 (t, J=6.9, 6H, CH<sub>3</sub>), 1.1-1.5 (m, 78H, CH<sub>2</sub>, CH<sub>3</sub>), 1.86 (t, J=6.6 Hz, 6H, CH<sub>3</sub>), 2.9-3.25 (m, 28H, CH<sub>2</sub>), 3.62 (s, 3H, CH<sub>3</sub>), 3.63 (s, 3H, CH<sub>3</sub>), 3.64 (s, 3H, CH<sub>3</sub>), 3.65 (s, 3H, CH<sub>3</sub>), 4.07 (q, J=7.5 Hz, 4H, CH<sub>2</sub>), 4.43 (t, J=7Hz, 4H, CH<sub>2</sub>), 10.07 (s, 3H, CH), 10.27 (s, 1H).

**Compound 102:** Trifluoroacetate (1 ml) was added to a solution of **101** (165 mg, 0.1 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2 ml) at 0<sup>0</sup> C, stirred for 1.5 h, and concentrated *in vacuo* at 20<sup>0</sup> C. The product was dried *in vacuo* for 3h, dissolved in 10 ml of 10% HCl, washed with CHCl<sub>3</sub>, and the aqueous part was concentrated and dried *in vacuo* at 20<sup>0</sup> C; yield 100 mg (75%), purple solid, mp >200<sup>0</sup> C. Purity 96% (HPLC). <sup>1</sup>H-NMR (D<sub>2</sub>O):  $\delta$  0.7-1.0 (m), 1.0-1.2 (m), 1.29 (t, J=7.3 Hz), 1.4-1.7 (m), 2.6-2.35 (m), 2.72 (t, J=7.2 Hz), 2.9-3.2 (m), 3.6-3.75 (m), 3.76 (s), 3.79 (s), 4.29 (q, J=7.6 Hz), 4.4-4.7 (m). MS-MALDI (m/z): 1193 (M<sup>+</sup> + 4 HCl), 1156 (M<sup>+</sup> + 3 HCl), 1119 (M<sup>+</sup> + 2 HCl), 1083 (M<sup>+</sup> + HCl), 1048 (M<sup>+</sup> + 1).

**Compound 103** was prepared from **71** in 85% yield as described above for **96**. <sup>1</sup>H-NMR (CDCl<sub>3</sub>):  $\delta$  0.97 (t, J=7 Hz, 3H, CH<sub>3</sub>), 1.3-1.55 (m, 16H, CH<sub>2</sub>), 2.27 (s, 3H, CH<sub>3</sub>), 2.29 (s, 9H, CH<sub>3</sub>), 2.55 (s, 24H, CH<sub>3</sub>), 3.0-3.25 (m, 16H, CH<sub>2</sub>), 3.31 (t, J=5.8 Hz, 2H, CH<sub>2</sub>), 4.41 (s, 2H, CH<sub>2</sub>), 6.90 (s, 2H, Ph), 6.92 (s, 6H, Ph).

**Compound 104** was prepared from **103** in 80% yield as described for **97**. <sup>1</sup>H-NMR (CDCl<sub>3</sub>):  $\delta$  1.10 (t, J=7.2 Hz, 3H, CH<sub>3</sub>), 1.5-1.8 (m, 16H, CH<sub>2</sub>), 2.5-2.75 (m, 16H, CH<sub>2</sub>), 3.57 (t, J=5.5 Hz, 2H, CH<sub>2</sub>). <sup>13</sup>C-NMR (CDCl<sub>3</sub>):  $\delta$  15.23, 27.55, 27.92, 28.58, 32.35, 44.02, 49.35, 49.66, 49.80, 62.32.

**Compound 105** was prepared from **104** in 98% yield as described for **98**. <sup>1</sup>H-NMR (CDCl<sub>3</sub>):  $\delta$  1.09 (t, J=7.1 Hz, 3H, CH<sub>3</sub>), 1.4-1.7 (m, 52 H, CH<sub>2</sub>, CH<sub>3</sub>), 3.05-3.3 (m, 16H, CH<sub>2</sub>), 3.67 (t, J=5.8 Hz, 2H, CH<sub>2</sub>).

**Compound 106:** Oxalyl chloride (2N solution in  $\text{CH}_2\text{Cl}_2$ , 0.821  $\mu\text{l}$ , 1.642 mmol) was diluted with anhydrous  $\text{CH}_2\text{Cl}_2$  (6 ml) at  $-60^\circ\text{C}$ . DMSO (223  $\mu\text{l}$ , 2.59 mmol) in  $\text{CH}_2\text{Cl}_2$  (3 ml) was added to the mixture, the latter stirred for 5 min at  $-60^\circ\text{C}$ , and **105** (1.115 g, 1.525 mmol) in  $\text{CH}_2\text{Cl}_2$  (9 ml) was introduced into the reaction. After 30 min of stirring at  $-60^\circ\text{C}$  triethylamine (1.06 ml, 14.46 mmol) was added into the reaction mixture and the temperature was allowed to rise to  $20^\circ\text{C}$  (ca. 1.5 h). The reaction mixture was diluted with  $\text{CH}_2\text{Cl}_2$ , washed with  $\text{H}_2\text{O}$ ,  $\text{NaHCO}_3$ , and brine. The product was concentrated to dryness *in vacuo* and purified by column chromatography ( $\text{SiO}_2$ , EtOAc/Hexane= 3:7); yield 989 mg (89%).  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  1.09 (t,  $J=7.0$  Hz, 3H,  $\text{CH}_3$ ), 1.4-1.6 (m, 48H,  $\text{CH}_2$ ,  $\text{CH}_3$ ), 1.84 (m, 2H,  $\text{CH}_2$ ), 2.45 (t,  $J=6.8$ , 2H,  $\text{CH}_2$ ), 3.05-3.3 (m, 16H,  $\text{CH}_2$ ), 9.78 (s, 1H, CHO).

**Compound 107:** Platinum oxide (100 mg) was reduced in methanol (30 ml) in the hydrogen atmosphere at 30 psi. for 15 min. Product **106** (989 mg, 1.36 mmol) was dissolved in a 2M solution of ethylamine in EtOH (7 ml), added to the hydrogenation flask, and hydrogenated for 10 h at 50 psi. The catalyst was removed by filtration through Celite and the filtrate was concentrated to dryness *in vacuo*; yield 1.0 g (99%).  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  1.09 (t,  $J=7.6$  Hz, 3H,  $\text{CH}_3$ ), 1.12 (t,  $J=7.2$  Hz, 3H,  $\text{CH}_3$ ), 1.3-1.65 (m, 50H,  $\text{CH}_2$ ,  $\text{CH}_3$ ), 1.66 (m, 2H,  $\text{CH}_2$ ), 2.71 (m, 2H,  $\text{CH}_2$ ), 3.1-3.3 (m, 18H,  $\text{CH}_2$ ). MS-MALDI ( $m/z$ ): 758.8 ( $\text{M}^+$ , 100%), 744 (30%).

**Compound 108** was prepared from **107** in 98% yield as described above for **101**.  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ):  $\delta$  0.8-1.8 (m), 1.87 (t,  $J=7.3$  Hz, 6H,  $\text{CH}_3$ ), 2.5-3.5 (m), 3.65 (s, 6H,  $\text{CH}_3$ ), 3.67 (s, 6H,  $\text{CH}_3$ ), 4.10 (q,  $J=7.5$ , 4H,  $\text{CH}_2$ ), 4.46 (t,  $J=6.8$ , 4H,  $\text{CH}_2$ ), 10.11 (bs, 4H, porph. core).

**Compound 109** was prepared from **108** in 75% yield as described above for **102**; purple solid, mp  $>200^\circ\text{C}$ . Purity 96% (HPLC).  $^1\text{H-NMR}$  ( $\text{D}_2\text{O}$ ):  $\delta$  0.25-0.5 (m), 0.6-0.8 (m), 1.1-1.5 (m), 1.5-2.0 (m), 1.32 (t), 2.6-3.4 (m), 3.6-3.9 (m), 3.78 (s), 3.82 (s), 4.2-4.6 (m). MS-MALDI ( $m/z$ ): 1246.22 ( $\text{M}^+ + 1$ ), 623.82 ( $\text{M}^{2+}$ ).

**Compound 111:** Nitrile **70** (274 mg, 0.343 mmol) was hydrogenated in a 5% solution of  $\text{CHCl}_3$  in EtOH (20 ml) in the presence of  $\text{PtO}_2$  (60 mg) at 50 psi for 15 h. The reaction mixture was filtered through Celite and concentrated to dryness *in vacuo* to yield **110** (288 mg, 100%). Amine **110** was utilized without further purification. Mesoporphyrin IX dihydrochloride (100 mg, 0.156 mmol) was dissolved in DMF (4 ml), HBTU (118 mg, 0.312













mmol) and DIEA (174  $\mu$ l, 2 mmol) were added to the solution, the latter stirred for 5 min, combined with **110**, and kept for 15 h. The reaction mixture was quenched with 0.5 ml of H<sub>2</sub>O, concentrated *in vacuo*, dissolved in CHCl<sub>3</sub>, washed with 3% HCl solution, brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and purified on a column (SiO<sub>2</sub>, CHCl<sub>3</sub>/MeOH=15:4); yield 327 mg (98%). <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  0.5-0.8 (m), 0.8-1.1 (m), 0.87 (t, J=7 Hz), 1.1-1.3 (m), 1.5-1.8 (m), 1.88 (q, J=5 Hz), 2.0-2.3 (m), 2.10 (s, CH<sub>3</sub>-Mes), 2.13 (s, CH<sub>3</sub>-Mes), 2.16 (s, CH<sub>3</sub>-Mes), 2.18 (s, CH<sub>3</sub>-Mes), 2.43 (s, CH<sub>3</sub>-Mes), 2.49 (s, CH<sub>3</sub>-Mes), 2.7-2.9 (m), 2.9-3.2 (m), 3.64 (s, porph), 3.66 (s, porph), 4.1 (q, porph.), 4.4 (t, porph.), 6.69 (s, NH), 6.80 (s, NH), 6.85 (s, NH), 6.93 (s, NH), 10.07 (s, porph), 10.09 (s, porph), 10.22 (s, porph.). MALDI (m/z): 2141.03 (M<sup>+</sup> +1), 1071.22 (M<sup>2+</sup>).











**Compound 112:** Porphyrin **111** (350 mg, 0.163 mmol) was stirred in a suspension of LiAlH<sub>4</sub> (12.4 mg, 0.326 mmol) in THF (1 ml) for 24 h at 50<sup>0</sup> C. The reaction mixture was cooled to 0<sup>0</sup> C, quenched with H<sub>2</sub>O (0.1 ml), basified with 2N NaOH (0.5 ml), then diluted with CH<sub>2</sub>Cl<sub>2</sub>, filtered through Celite, washed with H<sub>2</sub>O, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated *in vacuo*. The product was stirred in a mixture of phenol (920 mg, 9.4 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (8 ml), and 30% solution of HBr in AcOH (7 ml) for 15 h. The reaction mixture was cooled to 0<sup>0</sup> C, diluted with 10 ml of H<sub>2</sub>O and the aqueous layer was filtered and washed with CH<sub>2</sub>Cl<sub>2</sub>. Following drying *in vacuo* of the aqueous solution, the residue was basified with 2N NaOH (1 ml) and 16 N NaOH (1 ml) and the reaction product was extracted with CHCl<sub>3</sub>. The product **112** was purified by HPLC and converted to its hydrochloride by dissolution in 10% HCl and evaporation of the acid *in vacuo*; purple solid, mp 200<sup>0</sup> C. Purity 96% (HPLC). <sup>1</sup>H NMR (D<sub>2</sub>O):  $\delta$  1.32 (t, J=7.3 Hz, 6H, CH<sub>3</sub>), 1.25-1.35 (m, 30H, CH<sub>2</sub>), 2.5-2.65 (m, 4H, CH<sub>2</sub>), 3.0-3.3 (m, 32H, CH<sub>2</sub>), 3.4-3.5 (m, 4H, CH<sub>2</sub>), 3.55-3.65 (m, 6H, CH<sub>3</sub>), 3.73 (s, 3H, CH<sub>3</sub>), 3.75 (s, 3H, CH<sub>3</sub>), 4.0-4.25 (m, 4H, CH<sub>2</sub>), 4.3-4.5 (m, 4H). MALDI (m/z): 1073.8 (M<sup>+</sup> + NH<sub>4</sub>Cl), 1020.0 (M<sup>+</sup> + 1), 510.62 (M<sup>2+</sup>), 340.82 (M<sup>3+</sup>).

**Compound 113:** Diisobutylaluminum hydrate (1.16 ml of 1.5 M solution in toluene, 1.74 mmol) was added into a solution of mesoporphyrin IX dimethyl ester (500 mg, 84 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 ml) at -78<sup>0</sup> C, stirred at this temperature for 1h, quenched with saturated solution of NH<sub>4</sub>Cl (1 ml), followed by 3.7% solution of HCl (2 ml). The temperature of the reaction mixture was allowed to rise to 20<sup>0</sup>C, the product was extracted with CH<sub>2</sub>Cl<sub>2</sub>, dried (Na<sub>2</sub>SO<sub>4</sub>), and purified on a column (SiO<sub>2</sub>, EtOAc/Hexane=3:7), yield 330 mg (73%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>):  $\delta$  1.86 (t, J=7.6 Hz, 6H, CH<sub>3</sub>), 3.39 (t, J=7.4 Hz, 6H, CH<sub>3</sub>), 3.60 (s, 6H,

CH<sub>3</sub>), 3.62 (s, 6H, CH<sub>3</sub>), 4.0-4.2 (m, 4H, CH<sub>2</sub>), 4.25-4.45 (m, 4H, CH<sub>2</sub>), 9.97 (s, 1H), 10.04 (s, 1H), 10.05 (s, 1H), 10.058 (s, 1H), 10.062 (s, 1H), 10.07 (s, 1H).

Table 1. Polyamine analogs

Compound	Structure
SL-11090	 <chem>CNCCCCN[C@H]1C[C@H]2[C@@H]1CNCCCCN2</chem> $\cdot 4\text{HCl}$
SL-11091	 $\cdot 4\text{HCl}$
SL-11092	 $\cdot 4\text{HCl}$
SL-11101	 $\cdot 4\text{HCl}$
SL-11103	 $\cdot 4\text{HCl}$
SL-11114	 $4\text{HCl}$
SL-11118	 $4\text{HCl}$
SL-11121	 $5\text{HCl}$
SL-11122	 $5\text{HCl}$
SL-11123	 $5\text{HCl}$

SL-11124	 4HCl
SL-11126	 5 HCl
SL-11127	 5 HCl
SL-11128	 5 HCl
SL-11129	 5 HCl
SL-11130	 5 HCl
SL-11132	 4HCl
SL-11133	 5 HCl
SL-11134	 4HCl
SL-11136	 4HCl













SL-11137	 <p>4 HCl</p>
SL-11141	 <p>5 HCl</p>
SL-11144	 <p>10 HCl</p>
SL-11150	 <p>10 HCl</p>
SL-11201	 <p>10 HCl</p>
SL-11202	 <p>10 HCl</p>

Table 2. Polyamine-porphyrin conjugates

SL-11161	
SL-11162	
SL-11164	
SL-11177	
SL-11184	
SL-11202	

5

## EXAMPLE 2

***In vitro* testing of the efficacy of novel polyamine analogs against tumor cell lines**

These experiments are designed to evaluate the newly synthesized compounds described above against cultured human carcinoma cell lines for their effects on cell growth, cell-cycle regulation and polyamine regulatory responses. Additional methods for testing compounds are described in U.S. Patent No. 5,889,061.

10

As shown in Table 3 and Figs. 1-32, several novel conformationally restricted polyamine analogs were tested for anti-proliferative properties against cancer cells. Table 3 illustrates the concentration in  $\mu\text{M}$  of the various novel polyamine analogs needed for 50% growth inhibition ( $\text{ID}_{50}$ ) values for human cancer cell lines LNCaP, PC-3, DuPro (all three human prostate cancer cell lines), HT-29 (colon cancer cell line), A549 (lung cancer cell line), MCF7 (breast cancer cell line), and U251 MG-NCI (brain cancer cell line). Figs. 1-32 show a representative plot of the effects of some of these novel analogs on the growth of human tumor cell lines, as determined by MTT (methyl thiazol tetrazolium) assay; known anti-proliferative polyamine analogs BE-333, BE-343, BE-444, and BE-4444 were used for comparative purposes.

#### *Cell Lines and Media*

Human breast cancer cell line MCF7 was grown in Richter's Improved Modified Eagle's Medium supplemented with 10% fetal bovine serum (FBS) and 2.2 g/L sodium bicarbonate. Human brain tumor cell line U251 MG-NCI was grown in Dulbecco's Modified Eagle's Medium supplemented with 10% FBS. Human lung cancer cell line A549 was grown in Ham's F-12K medium (Cellgro, Mediatech, Inc., VA), supplemented with 10% FBS and 2 mM L-glutamine. Human colon cancer cell line HT29 was cultured in McCoy's 5A medium (Gibco, BRL, Gaithersburg, MD) supplemented with 10% FBS. Human prostate cancer cell lines PC-3, LNCAP and DuPro were grown in RPMI 1640 Medium (Cellgro, Mediatech, Inc., VA) supplemented with 10% FBS. Another human prostate cancer cell line DU145 was grown in Dulbecco's Modified Eagle's Medium (Gibco, BRL, Gaithersburg, MD) supplemented with 5% FBS. The A549, MCF7, PC3, LNCAP and DuPro cell lines were cultured in 100 units/mL penicillin and 100  $\mu\text{g/mL}$  streptomycin. HT29 and U251MG cell lines were grown in 50  $\mu\text{g/mL}$  gentamycin (Gibco, BRL, Gaithersburg, MD). DU145 cell line was maintained in 1% antibiotic-antimycotic solution (Sigma, St. Louis, MO). The cell cultures were maintained at 37°C in 5%CO<sub>2</sub>/95% humidified air. DuPro cells were obtained from M. Eileen Dolan, University of Chicago. All other cells are available from the American Type Culture Collection, Rockville, MD.

#### *MTT assay*

A conventional MTT assay was used to evaluate percent cell survival.

Exponentially growing monolayer cells were plated in 96-well plates at a density of 500 cells per well and allowed to grow for 24 hours. Serial dilutions of the drugs were added to the wells. Six days after drug treatment, 25  $\mu$ l of MTT solution (5 mg/ml) was added to each well and incubated for 4 hours at 37°C. Then 100  $\mu$ l of lysis buffer (20% sodium dodecyl sulfate, 50% DMF, and 0.8% acetic acid, pH 4.7) was added to each well and incubated for an additional 22 hours. A microplate reader ("EMAX"-brand, Molecular Devices, Sunnyvale, Calif.) set at 570 nm was used to determine the optical density of the cultures. Results are expressed as a ratio of the optical density in drug-treated wells to the optical density in wells treated with vehicle only.

**TABLE 3**

	PC-3	DU-145	DUPRO	HT-29	A549	MCF7	U251M G
BE-4444	0.54	0.07	0.2	0.8	0.4	>31.25	NT
SL-11029	24.5	0.32	NT	>31.25	>31.25	>31.25	>31.25
SL-11090	>31.25	>31.25	NT	>31.25	>31.25	>31.25	>31.25
SL-11091	>31.25	1.33	NT	>31.25	>31.25	>31.25	>31.25
SL-11092	>31.25	1.7	NT	>31.25	>31.25	>31.25	>31.25
SL-11093	14.3	0.01	0.06	0.40	0.26	0.66	NT
SL-11094	>31.25	12.6	NT	28.8	>31.25	>31.25	>31.25
SL-11098	1.4	0.018	0.08	0.40	0.51	>31.25	0.10
SL-11099	2.5	0.014	0.08	1.00	0.65	26.3	0.11
SL-11100	4.7	0.021	0.29	2.00	2.20	>31.25	0.22
SL-11101	7.7	0.218	0.85	5.20	0.15	>31.25	1.70
SL-11102	>31.25	0.027	0.15	0.73	12.40	>31.25	0.15
SL-11103	>31.25	2.8	NT	29.4	>31.25	>31.25	9.50
SL-	>31.25	9.4	NT	25.8	0.43	>31.25	14.71



11104							
SL-11105	>31.25	1.6	>31.25	25.2	>31.25	>31.25	25.9
SL-11108	2.2	0.13	0.98	2.00	>31.25	>31.25	2.00
SL-11114	0.70	0.135	0.64	3.6	>31.25	NT	NT
SL-11118	1.65	0.05	0.25	0.98	0.21	NT	NT
SL-11119	>31.25	0.08	0.44	0.97	NT	NT	NT
SL-11121	0.52	0.08	0.40	0.80	>31.25	17.0	NT
SL-11122	>31.25	0.80	0.56	0.80	>31.25	>31.25	NT
SL-11123	>31.25	0.51	>31.25	10.42	>31.25	>31.25	NT
SL-11124	>31.25	>31.25	>31.25	>31.25	>31.25	>31.25	NT
SL-11126	0.20	0.51	1.10	1.50	>31.25	0.70	NT
SL-11127	>31.25	0.22	1.3	2.91	NT	NT	NT
SL-11128	0.50	0.14	1.25	1.35	NT	NT	NT
SL-11129	1.70	0.32	NT	NT	NT	NT	NT
SL-11130	>31.25	0.43	NT	NT	NT	NT	NT

NT indicates not tested.

5

### EXAMPLE 3

#### *In vivo* testing of anti-tumor activity of polyamine analogs

10      Analogs found to have potent or mechanism-based anti-proliferative activity *in vitro* towards cultured carcinoma cells are evaluated in *in vivo* model systems. The first goal is to determine the relative toxicity of the analogs in non-tumor-bearing animals, such as DBA/2 mice. Groups of three animals each are injected intraperitoneally with increasing concentrations of an analog, beginning at 10 mg/kg. Toxicity, as indicated by morbidity, is

5 closely monitored over the first 24 hr. A well-characterized polyamine analog, such as BE-333, is used as an internal standard in these studies, since a data base has already been established regarding acute toxicity via a single dose treatment relative to chronic toxicity via a daily x 5 d schedule. Thus, in the case of new analogs, single dose toxicity relative to BE-333 is used to project the range of doses to be used on a daily x 5 d schedule.

10 After the highest tolerated dosage on a daily x 5 d schedule is deduced, antitumor activity is determined. Tumors are subcutaneously implanted into nude athymic mice by trocar and allowed to reach 100-200 mm<sup>3</sup> before initiating treatment by intraperitoneal injection daily x 5 d. Analogs are given in a range between 10 and 200 mg/kg. Analogs are evaluated at three treatment dosages with 10-15 animals per group (a minimum of three from each are used for pharmacodynamic studies, described below). Mice are monitored and weighed twice weekly to determine tumor size and toxicity. Tumor size is determined by multi-directional measurement from which volume in mm<sup>3</sup> is calculated. Tumors are followed until median tumor volume of each group reaches 1500 mm<sup>3</sup> (i.e., 20% of body weight), at which time the animals are sacrificed. Although the initial anti-tumor studies focused on a daily x 5 d schedule, constant infusion is performed via Alzet pump delivery for 5 days since this schedule dramatically improves the anti-tumor activity of BE-333 against A549 human large cell lung carcinoma. Sharma et al. (1997) *Clin. Cancer Res.* 3:1239-1244. In addition to assessing anti-tumor activity, free analog levels in tumor and normal tissues are determined in test animals.

#### EXAMPLE 4

##### *In vitro testing of porphyrin-polyamine conjugates*

25 Using the protocols of Example 2, the porphyrin-polyamine conjugates were tested for activity *in vitro* in various cancer cell lines. The results are shown in Table 4 and Figs. 45-49.

**Table 4**

**Effect of Polyamine Porphyrin Complexes on the Growth of Human Prostate Cell Lines by MTT Assay.**

Compounds	IC <sub>50</sub> (μM) in different cell lines			
	PC3	DUPRO	LNCAP	DU145
SL-11162	2.8	1.9	4.8	
SL-11177	0.9	0.7	4.8	
SL-11184	1.4	1.5		8.4
SL-11202	1.15	1.3		4.6

## EXAMPLE 5

### *In vivo testing of porphyrin-polyamine conjugates*

5 Conjugates found to have potent or mechanism-based anti-proliferative activity *in vitro* towards cultured carcinoma cells are evaluated in *in vivo* model systems. The first goal is to determine the relative toxicity of the conjugates in non-tumor-bearing animals, such as DBA/2 mice. Groups of three animals each are injected intraperitoneally with increasing concentrations of an conjugate, beginning at 10 mg/kg. Toxicity, as indicated by morbidity, is closely monitored over the first 24 hr. A well-characterized compound, such as BE-333, is used as an internal standard in these studies, since a data base has already been established regarding acute toxicity via a single dose treatment relative to chronic toxicity via a daily x 5 d schedule. Thus, in the case of new conjugates, single dose toxicity relative to BE-333 is used to project the range of doses to be used on a daily x 5 d schedule.

15 After the highest tolerated dosage on a daily x 5 d schedule is deduced, antitumor activity is determined. Tumors are subcutaneously implanted into nude athymic mice by trocar and allowed to reach 100-200 mm<sup>3</sup> before initiating treatment by intraperitoneal injection daily x 5 d. Conjugates are given in a range between 10 and 200 mg/kg. Conjugates are evaluated at three treatment dosages with 10-15 animals per group (a minimum of three from each are used for pharmacodynamic studies, described below). Mice are monitored and weighed twice weekly to determine tumor size and toxicity. Tumor size is determined by multi-directional measurement from which volume in mm<sup>3</sup> is calculated. Tumors are followed until median tumor volume of each group reaches 1500 mm<sup>3</sup> (i.e., 20% of body weight), at which time the animals are sacrificed. Although the initial anti-tumor studies focused on a daily x 5 d schedule, constant infusion is performed via Alzet pump delivery for 5 days since this schedule dramatically improves the anti-tumor activity of BE-333 against A549 human large cell lung carcinoma. Sharma et al.

(1997) *Clin. Cancer Res.* 3:1239-1244. In addition to assessing anti-tumor activity, free conjugate levels in tumor and normal tissues are determined in test animals.

5 Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it is apparent to those skilled in the art that certain minor changes and modifications will be practiced. Therefore, the description and examples should not be construed as limiting the scope of the invention, which is delineated by the appended claims. All references disclosed herein, including U.S. Patent No. 5,889,061, are hereby incorporated by reference in their entirety.

## CLAIMS

What is claimed is:

5

1. A conformationally restricted polyamine analog of the formula:

E-NH-B-A-B-NH-B-A-B-NH-B-A-B-NH-B-A-B-NH-E

wherein each A is independently selected from the group consisting of: a single bond,  
C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub>  
10 cycloalkenyl;

each B is independently selected from the group consisting of: a single bond, C<sub>1</sub>-C<sub>6</sub> alkyl,  
and C<sub>2</sub>-C<sub>6</sub> alkenyl;


and each E is independently selected from the group consisting of H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub>  
alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl;

15


with the proviso that either at least one A moiety is selected from the group consisting of  
C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl,  
or at least one B moiety is selected from the group consisting of C<sub>2</sub>-C<sub>6</sub> alkenyl;  
and any salt or stereoisomer thereof.

20

2. A conformationally restricted polyamine analog according to claim 1, selected from  
the group consisting of



25




5 , and any salt or stereoisomer thereof.

3. A conformationally restricted polyamine analog of the formula:




- 10 wherein each A is independently selected from the group consisting of: a single bond, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl;
- each B is independently selected from the group consisting of: a single bond, C<sub>1</sub>-C<sub>6</sub> alkyl, and C<sub>2</sub>-C<sub>6</sub> alkenyl;
- 15 each E is independently selected from the group consisting of H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl; and x is an integer from 2 to 16;
- with the proviso that either at least one A moiety is selected from the group consisting of C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl,
- 20 or at least one B moiety is selected from the group consisting of C<sub>2</sub>-C<sub>6</sub> alkenyl; and any salt or stereoisomer thereof.

4. A conformationally restricted polyamine analog according to claim 3, selected from the group consisting of:



and




;

and any salt or stereoisomer thereof.

5

5. A polyamine analog of the formula:



wherein each A is independently selected from the group consisting of: a single bond, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl;  
each B is independently selected from the group consisting of: a single bond, C<sub>1</sub>-C<sub>6</sub> alkyl, and C<sub>2</sub>-C<sub>6</sub> alkenyl;

10

each E is independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkanol, C<sub>3</sub>-C<sub>6</sub> cycloalkanol, and C<sub>3</sub>-C<sub>6</sub> hydroxyaryl,

with the proviso that at least one E moiety be selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkanol, C<sub>3</sub>-C<sub>6</sub> cycloalkanol, and C<sub>3</sub>-C<sub>6</sub> hydroxyaryl;

15


and x is an integer from 0 to 16;


and any salt or stereoisomer thereof.

6. A polyamine analog according to claim 5, selected from the group consisting


20

of:





, and



;

and all salts and stereoisomers thereof.

5

7. A conformationally restricted polyamine analog of the formula:

E-NH-D-NH-B-A-B-NH-D-NH-E

wherein each A is selected from the group consisting of C<sub>2</sub>-C<sub>6</sub> alkynyl;

each B is independently selected from the group consisting of: a single bond, C<sub>1</sub>-C<sub>6</sub> alkyl,

10 and C<sub>2</sub>-C<sub>6</sub> alkenyl;

each D is independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl,

C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloalkenyl, and C<sub>3</sub>-C<sub>6</sub> cycloaryl; and

each E is independently selected from the group consisting of H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub>

alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl;


15 and any salt or stereoisomer thereof.

8. A conformationally restricted polyamine analog according to claim 7, selected from the group consisting of

20




,




, and





and any salt or stereoisomer thereof.

5            9. A conformationally restricted polyamine analog of the formula:



wherein F is selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl;



each A is independently selected from the group consisting of: a single bond, C<sub>1</sub>-C<sub>6</sub> alkyl; C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl;

10        each B is independently selected from the group consisting of: a single bond, C<sub>1</sub>-C<sub>6</sub> alkyl, and C<sub>2</sub>-C<sub>6</sub> alkenyl; and

each E is independently selected from the group consisting of H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl;


15        with the proviso that either at least one A moiety is selected from the group consisting of C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl, or at least one B moiety is selected from the group consisting of C<sub>2</sub>-C<sub>6</sub> alkenyl; and any salt or stereoisomer thereof.

20        10. A conformationally restricted polyamine analog according to claim 9, selected from the group consisting of



25        and any salt or stereoisomer thereof.

11. A conformationally restricted polyamine analog of the formula:





wherein A is independently selected from the group consisting of: a single bond, C<sub>1</sub>-C<sub>6</sub> alkyl; C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl;

each B is independently selected from the group consisting of: a single bond, C<sub>1</sub>-C<sub>6</sub> alkyl, and C<sub>2</sub>-C<sub>6</sub> alkenyl;


with the proviso that A and both B moieties are not all a single bond;

and any salt or stereoisomer thereof.

12. A conformationally restricted polyamine analog according to claim 11, selected from the group consisting of:



, and



and any salt or stereoisomer thereof.

5                    13. A conformationally restricted polyamine analog of the formula:

E-NH-D-NH-B-A-B-NH-D-NH-E

wherein A is selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloalkenyl, and C<sub>3</sub>-C<sub>6</sub> cycloaryl;

each B is independently selected from the group consisting of: a single bond, C<sub>1</sub>-C<sub>6</sub> alkyl,  
10                    and C<sub>2</sub>-C<sub>6</sub> alkenyl;

each D is independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloalkenyl, and C<sub>3</sub>-C<sub>6</sub> cycloaryl; and

each E is independently selected from the group consisting of H and C<sub>1</sub>-C<sub>6</sub> alkyl;

with the proviso that either at least one A moiety is selected from the group consisting of  
15                    C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl,

or at least one B moiety is selected from the group consisting of C<sub>2</sub>-C<sub>6</sub> alkenyl;

and the proviso that at least one E is selected from the group consisting of H or methyl;


and any salt or stereoisomer thereof.

20                    14. A conformationally restricted polyamine analog according to claim 13 of the formula



and any salt or stereoisomer thereof.

15. A conformationally restricted polyamine analog of the formula:



wherein A is *trans*-ethene

each B is independently selected from the group consisting of: a single bond, C<sub>1</sub>-C<sub>6</sub> alkyl, and C<sub>2</sub>-C<sub>6</sub> alkenyl;

each D is independently selected from the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl, with the proviso that D is not C<sub>3</sub> alkyl; and

each E is independently selected from the group consisting of H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>6</sub> cycloalkyl, C<sub>3</sub>-C<sub>6</sub> cycloaryl, and C<sub>3</sub>-C<sub>6</sub> cycloalkenyl; and any salt thereof.

16. The polyamine analog of claims 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15, further comprising a pharmaceutically acceptable excipient.

17. A method of treating an indication in an individual comprising the step of administering to the individual a therapeutic amount of a polyamine analog of claims 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15.

18. The method of claim 17, wherein the individual is a human.

19. The method of claim 17, wherein the indication is a cancer.

20. The method of claim 19, wherein the cancer affects cells of the bladder, blood, brain, breast, colon, digestive tract, lung, ovaries, pancreas, prostate gland, or skin.

21. The method of claim 17, wherein the indication is Alzheimer's disease, epilepsy, multiple sclerosis, problems associated with tissue grafts and organ transplants, psoriasis, restenosis, stomach ulcers, or tissue overgrowth after surgery, or an infection or infestation of parasites, bacteria, fungi or insects.

22. A method of suppressing cell growth in an individual comprising the step of administering to the individual a therapeutic amount of a polyamine analog of claims 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15.

5


23. The method of claim 22, wherein the individual is a human.

24. A porphyrin-polyamine conjugate, comprising a polyamine compound covalently linked to a porphyrin compound.

10

25. A compound of claim 24, wherein the covalent link is an amide bond or a amine bond.

26. A compound according to claim 24 of the formula



15

wherein at least one of  $J_1$ - $J_8$  is independently selected from the group consisting of  $-K_1-G-L-(N(P)-A)_n-K_2$

where  $K_1$  is independently selected from the group consisting of  $C_1$ - $C_8$  alkyl and where the valence to the left of  $K_1$  attaches to the porphyrin ring;

20  $G$  is  $-O-$ ,  $-(C=O)-$ ,  $-C(=O)-O-$ ,  $-O-(C=O)-$ ,  $-O-(C=O)-O-$ ,  $-O-(C=O)-N-$ ,  $-N-(C=O)-O-$ , or a nonentity;

L is C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>3</sub>-C<sub>8</sub> cycloaryl, C<sub>1</sub>-C<sub>8</sub> alkoxy, C<sub>1</sub>-C<sub>8</sub> alkyl-C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>1</sub>-C<sub>8</sub> alkyl-C<sub>3</sub>-C<sub>8</sub> cycloaryl, C<sub>1</sub>-C<sub>8</sub> alkoxy-C<sub>3</sub>-C<sub>8</sub> cycloaryl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl-C<sub>3</sub>-C<sub>8</sub> cycloaryl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl-C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>3</sub>-C<sub>8</sub> cycloaryl-C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>3</sub>-C<sub>8</sub> cycloaryl-C<sub>1</sub>-C<sub>8</sub> alkoxy, C<sub>3</sub>-C<sub>8</sub> cycloaryl-C<sub>3</sub>-C<sub>8</sub> cycloalkyl, or a nonentity;

5 each A is independently selected from the group consisting of C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>2</sub>-C<sub>8</sub> alkenyl, C<sub>2</sub>-C<sub>8</sub> alkynyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>3</sub>-C<sub>8</sub> cycloaryl, C<sub>3</sub>-C<sub>8</sub> cycloalkenyl, and C<sub>3</sub>-C<sub>8</sub> cycloalkynyl;

P is selected from the group consisting of H and C<sub>1</sub>-C<sub>8</sub> alkyl;

n is an integer from 2 to 8;

10 and K<sub>2</sub> is independently selected from the group consisting of H, C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>2</sub>-C<sub>8</sub> alkenyl, C<sub>2</sub>-C<sub>8</sub> alkynyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>3</sub>-C<sub>8</sub> cycloaryl, C<sub>3</sub>-C<sub>8</sub> cycloalkenyl, C<sub>3</sub>-C<sub>8</sub> cycloalkynyl, C<sub>1</sub>-C<sub>8</sub> alkanol, C<sub>3</sub>-C<sub>8</sub> cycloalkanol, and C<sub>3</sub>-C<sub>8</sub> hydroxyaryl;

where the remainder of the groups J<sub>1</sub>-J<sub>8</sub> are each independently selected from the group consisting of H, K<sub>3</sub>, or K<sub>4</sub>-COOH, where K<sub>3</sub> is independently selected from the group  
15 consisting of C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>2</sub>-C<sub>8</sub> alkenyl, C<sub>2</sub>-C<sub>8</sub> alkynyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>3</sub>-C<sub>8</sub> cycloaryl, C<sub>3</sub>-C<sub>8</sub> cycloalkenyl, C<sub>3</sub>-C<sub>8</sub> cycloalkynyl, C<sub>1</sub>-C<sub>8</sub> alkanol, C<sub>3</sub>-C<sub>8</sub> cycloalkanol, and C<sub>3</sub>-C<sub>8</sub> hydroxyaryl; and K<sub>4</sub> is independently selected from the group consisting of C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>2</sub>-C<sub>8</sub> alkenyl, C<sub>2</sub>-C<sub>8</sub> alkynyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>3</sub>-C<sub>8</sub> cycloaryl, C<sub>3</sub>-C<sub>8</sub> cycloalkenyl, and C<sub>3</sub>-C<sub>8</sub> cycloalkynyl;

20 and any salt or stereoisomer thereof.

27. A compound according to claim 26, wherein K<sub>1</sub> and each A are independently selected from the group consisting of C<sub>1</sub>-C<sub>8</sub> alkyl, K<sub>2</sub> is independently selected from the group consisting of H and C<sub>1</sub>-C<sub>8</sub> alkyl, and the remainder of the groups J<sub>1</sub>-J<sub>8</sub> are each  
25 independently selected from the group consisting of H, K<sub>3</sub>, or K<sub>4</sub>-COOH, where K<sub>3</sub> and K<sub>4</sub> are independently selected from the group consisting of C<sub>1</sub>-C<sub>8</sub> alkyl.

28. A compound according to claim 27, wherein K<sub>1</sub> is -CH<sub>2</sub>-CH<sub>2</sub>- or -CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-.

30 29. A compound according to claim 27, wherein n is 4.

30. A compound according to claim 27, wherein J<sub>3</sub>, J<sub>4</sub>, J<sub>7</sub> and J<sub>8</sub> are independently selected from the group consisting of C<sub>1</sub>-C<sub>3</sub> alkyl;  
J<sub>5</sub> and J<sub>6</sub> are independently selected from the group consisting of C<sub>1</sub>-C<sub>3</sub> alkyl and C<sub>1</sub>-C<sub>3</sub> alkyl-COOH;

5 and J<sub>1</sub> and J<sub>2</sub> are independently selected from the group consisting of C<sub>1</sub>-C<sub>3</sub> alkyl-G-N(P<sub>1</sub>)-A-(NH-A)<sub>n1</sub>-K<sub>5</sub>;

wherein G is -(C=O)- or a nonentity;

P<sub>1</sub> is H or C<sub>1</sub>-C<sub>3</sub> alkyl;


each A is independently selected from the group consisting of C<sub>1</sub>-C<sub>8</sub> alkyl;

10 n<sub>1</sub> is 3 or 4;

and K<sub>5</sub> is independently selected from the group consisting of H and C<sub>1</sub>-C<sub>8</sub> alkyl.

31. A method of treating an indication in an individual comprising the step of administering to the individual a therapeutic amount of a porphyrin-polyamine conjugate of  
15 claim 24.

1/56





2/56




FIG. 2




FIG. 3





FIG. 4

5/56



6/56




FIG. 6

7/56




FIG. 7

8/56




FIG. 8

9/56




FIG. 9



10/56




FIG. 10

11/56




FIG. 11

12/56




FIG. 12

13/56




FIG. 13

14/56




FIG. 14

15/56




FIG. 15A

16/56




FIG. 15B

17/56




FIG. 16 A



18/56




FIG. 16 B

19/56




FIG. 17 A

20/56




FIG. 17 B

21/56




FIG. 18 A




FIG. 18 B




FIG. 19A

24/56




FIG. 19B




FIG. 20 A






FIG. 20 B




FIG. 21 A




FIG. 21 B




FIG. 22 A




FIG. 22 B




FIG. 23




FIG. 24




FIG. 25






FIG. 26




FIG. 27








FIG. 28











Figure 34








Figure 35










Figure 36






Figure 37








Figure 38



SCHEME 22

Figure 39



SCHEME 23




Figure 40A

SCHEME 24




Figure 40B

## SCHEME 24 (continued)




Figure 41

SCHEME 25










FIGURE 43


SCHEME 27





## FIGURE 44


## SCHEME 28














(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
9 November 2000 (09.11.2000)

PCT

(10) International Publication Number  
**WO 00/66587 A3**

(51) International Patent Classification<sup>7</sup>: **C07D 487/22**,  
C07C 211/22, 211/23, 211/18, 215/14, 257/18, A61P  
35/00, A61K 31/155, 31/13, 31/40

**BASU, Hirak, S.** [US/US]; 7226 Colony Drive, Madison,  
WI 53717 (US).

(21) International Application Number: **PCT/US00/11591**

(74) Agents: **CERPA, Robert, K. et al.**; Morrison & Foerster  
LLP, 755 Page Mill Road, Palo Alto, CA 94304-1018 (US).

(22) International Filing Date: **27 April 2000 (27.04.2000)**

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE,  
DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,  
ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,  
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ,  
PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT,  
TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:  
60/131,779 30 April 1999 (30.04.1999) **US**

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent  
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent  
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,  
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

Published:

(75) Inventors/Applicants (*for US only*): **FRYDMAN, Ben-**  
**jamin** [AR/US]; 821 N. Holt Circle, Madison, WI 53719  
(US). **MARTON, Laurence, J.** [US/US]; 5810 Tree Line  
Drive, Madison, WI 53711 (US). **REDDY, Venodhar, K.**  
[IN/US]; 4446 Hillcrest Drive #3, Madison, WI 53705  
(US). **VALASINAS, Aldonia** [AR/US]; 821 N. Holt  
Circle, Madison, WI 53719 (US). **BLOKHIN, Andrei, V.**  
[RU/US]; 14 Dorchester Circle, Madison, WI 53719 (US).

— *With international search report.*

(88) Date of publication of the international search report:  
25 January 2001

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: **POLYAMINES AND THEIR USE IN THERAPY**

(57) Abstract: Novel conformationally restricted polyamine analogs are provided, as well as compositions comprising these novel polyamine analogs. Methods of using the novel polyamine analogs in treatment of diseases such as cancer are also provided. Also provided is a method of delivering these analogs specifically to tumor cells by covalently attaching polyamine analogs to porphyrin compounds, along with novel polyamine-porphyrin covalent conjugates.

**WO 00/66587 A3**



## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/11591

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D487/22 C07C211/22 C07C211/23 C07C211/18 C07C215/14  
C07C257/18 A61P35/00 A61K31/155 A61K31/13 A61K31/40

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D C07C A61P A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, WPI Data, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	PATENT ABSTRACTS OF JAPAN vol. 1996, no. 02, 29 February 1996 (1996-02-29) & JP 07 277964 A (NIPPON KAYAKU CO LTD), 24 October 1995 (1995-10-24) abstract ---	1-4, 16-23
Y	WO 95 18091 A (CIBA GEIGY AG ;FREI JOERG (CH); STANEK JAROSLAV (CH)) 6 July 1995 (1995-07-06) page 31 -page 49; examples 1-23 ---	1-4, 16-23
Y	US 5 541 230 A (BASU HIRAK S ET AL) 30 July 1996 (1996-07-30) column 6, line 30 -column 20, line 64 --- -/-	1-4, 16-23

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance  
 "E" earlier document but published on or after the international filing date  
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)  
 "O" document referring to an oral disclosure, use, exhibition or other means  
 "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention  
 "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone  
 "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.  
 "&" document member of the same patent family

Date of the actual completion of the international search

26 October 2000

Date of mailing of the international search report

08.11.2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Bader, K

## INTERNATIONAL SEARCH REPORT

In/ tional Application No

PCT/US 00/11591

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 02027 A (TEVA PHARMA ;GILAD GAD (IL); GILAD VARDA (IL); STERLING JEFFREY (I) 23 January 1997 (1997-01-23) page 44 ----	13,17,18
X	WO 96 22962 A (UNIV COLUMBIA ;NAKANISHI KOJI (US); HUANG DANWEN (US); CHOI SEOK K) 1 August 1996 (1996-08-01) page 19 ----	13,17,18
X	PATENT ABSTRACTS OF JAPAN vol. 017, no. 327 (C-1073), 22 June 1993 (1993-06-22) & JP 05 032902 A (IDEMITSU PETROCHEM CO LTD;OTHERS: 01), 9 February 1993 (1993-02-09) abstract ----	13
X	US 4 590 288 A (KLEMMANN LAWRENCE P) 20 May 1986 (1986-05-20) column 3 -column 6 ----	13
X	US 4 092 432 A (BJORKLUND CHRISTER JAMES AXEL ET AL) 30 May 1978 (1978-05-30) column 4 -column 5; examples 1-6 ----	13
X	DE 12 95 826 B (J.R.GEIGY AG, BASEL (SCHWEIZ)) column 3; tables 1,NR.7 ----	13
X	US 5 889 061 A (MARTON LAURENCE J ET AL) 30 March 1999 (1999-03-30) column 19 -column 20; table 1 column 39; claims 15-22 ----	13,16-23
X	US 5 021 571 A (SRIVASTAVA SURESH C ET AL) 4 June 1991 (1991-06-04) column 6 ----	13,16-23
X	WO 98 17624 A (WISCONSIN ALUMNI RES FOUND) 30 April 1998 (1998-04-30) page 29; table 1 ----	13,16-23
X	US 4 035 174 A (GRIER NATHANIEL ET AL) 12 July 1977 (1977-07-12) column 18, line 20 ----	13
X	US 4 153 567 A (KLUGER EDWARD W ET AL) 8 May 1979 (1979-05-08) column 3, line 55 - line 60 ----	13
P,X	US 5 962 533 A (BERGERON JR RAYMOND J) 5 October 1999 (1999-10-05) page 7 -page 14; tables 1,2 ----	13,16-23

-/--

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	FISCHER, H. A.: "Synthesis of 3H-spermine" J. LABELLED COMPD., vol. 11, no. 1, 1975, pages 141-143, XP000956379 page 141 -page 143 ----	7,13
X	REDDY, VENODHAR K. ET AL: "Conformationally Restricted Analogs of 1N,12N-Bisethylspermine: Synthesis and Growth Inhibitory Effects on Human Tumor Cell Lines" J. MED. CHEM., vol. 41, no. 24, 1998, pages 4723-4732, XP000952845 page 4724; figure 1 ----	7,13, 16-23
X	GOTO, MASAFUMI ET AL: "Stereochemical studies of metal chelates. III. Preparation and stereochemistry of cobalt(III) complexes with C-substituted triethylenetetramines at the central ethylenediamine bridge" INORG. CHEM., vol. 8, no. 2, 1969, pages 358-366, XP000952924 page 361; figure 1 ----	13
X	ALFONSO, IGNACIO ET AL: "Sequential biocatalytic resolution of (+-)-trans-cyclohexane-1,2- diamine. Chemoenzymic synthesis of an optically active polyamine" CHEM. COMMUN. (CAMBRIDGE), no. 21, 1996, pages 2471-2472, XP000952857 page 2472 ----	13
X	NAGARAJAN, SRINIVASAN ET AL: "Chemistry of naturally occurring polyamines. 11. Unsaturated spermidine and spermine derivatives" J. ORG. CHEM., vol. 52, no. 22, 1987, pages 5044-5046, XP000952844 Scheme 1 page 5044 ----	7,13
X	SNYDER, RONALD D. ET AL: "Effects of polyamine analogs on the extent and fidelity of in vitro polypeptide synthesis" BIOCHEM. BIOPHYS. RES. COMMUN., vol. 176, no. 3, 1991, pages 1383-1392, XP000952839 page 1385; table 1 ----	13
	----- -/--	

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/11591

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WUNZ, TIMOTHY P. ET AL: "New antitumor agents containing the anthracene nucleus" J. MED. CHEM., vol. 30, no. 8, 1987, pages 1313-1321, XP000867863 scheme 1 page 1314 -----	13
X	KOBIRO, KAZUYA ET AL: "Synthesis and molecular structures of nickel(II) alkyl-substituted cyclam complexes" INORG. CHEM., vol. 31, no. 4, 1992, pages 676-685, XP000957409 page 677 -----	13

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 00/11591

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
  
Although claims 17-23,31 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☒ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:  
  
1-4,7,8,11-14,16-31 (therof 16-23 and 31 only partially)
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-4 (completely), 16-23 (in part)

A compound according to the definition of claim 1.

2. Claims: 5,6 (completely), 16-23 (in part)

A compound according to the definition of claim 5.

3. Claims: 7,8,13,14 (completely), 16-23 (in part)

A compound according to definition of claim 7 or according to the definition of claim 13.

4. Claims: 9,10 (completely), 16-23 (in part)

A compound according to definition of claim 9.

5. Claims: 11,12 (completely), 16-23 (in part)

A compound according to the definition of claim 11.

6. Claims: 15; 16-23 (in part)

A compound according to the definition of claim 15.

7. Claims: 24-31

A porphyrin-polyamine conjugate and its use.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/11591

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
JP 07277964	A	24-10-1995	NONE	
WO 9518091	A	06-07-1995	AT 169614 T AU 690618 B AU 1274495 A BR 9405892 A CN 1121707 A DE 69412467 D DE 69412467 T DK 686143 T EP 0686143 A ES 2119371 T JP 8507548 T NZ 277503 A US 5627215 A	15-08-1998 30-04-1998 17-07-1995 12-12-1995 01-05-1996 17-09-1998 28-01-1999 17-05-1999 13-12-1995 01-10-1998 13-08-1996 27-07-1997 06-05-1997
US 5541230	A	30-07-1996	US 5880161 A	09-03-1999
WO 9702027	A	23-01-1997	US 5646188 A AU 6452596 A EP 0865276 A	08-07-1997 05-02-1997 23-09-1998
WO 9622962	A	01-08-1996	US 6001824 A AU 5168496 A	14-12-1999 14-08-1996
JP 05032902	A	09-02-1993	NONE	
US 4590288	A	20-05-1986	NONE	
US 4092432	A	30-05-1978	SE 417569 B AR 216638 A AU 505423 B AU 1855076 A BR 7606992 A CA 1072008 A CH 625208 A DD 128387 A DE 2647915 A DK 473376 A,B, FR 2328399 A GB 1570517 A HU 176545 B IL 50649 A IN 143379 A IT 1121707 B JP 1119495 C JP 52051021 A JP 57007605 B MX 4511 E NL 7611508 A,B, SE 7511852 A SU 852169 A ZA 7606127 A	30-03-1981 15-01-1980 22-11-1979 20-04-1978 06-09-1977 19-02-1980 15-09-1981 16-11-1977 05-05-1977 23-04-1977 20-05-1977 02-07-1980 28-03-1981 30-06-1980 12-11-1977 23-04-1986 28-10-1982 23-04-1977 12-02-1982 02-06-1982 26-04-1977 23-04-1977 30-07-1981 28-09-1977
DE 1295826	B		NONE	
US 5889061	A	30-03-1999	NONE	

# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 00/11591

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5021571 A	04-06-1991	US 5089663 A US 5334729 A	18-02-1992 02-08-1994
WO 9817624 A	30-04-1998	AU 4900797 A EP 0934249 A	15-05-1998 11-08-1999
US 4035174 A	12-07-1977	US 4049559 A	20-09-1977
US 4153567 A	08-05-1979	NONE	
US 5962533 A	05-10-1999	NONE	